arXiv:math-ph/0312066v1 24 Dec 2003

Spectral asymptotics of harmonic oscillator perturbed by bounded potential

Markus Klein [*] and Evgeni Korotyaev [†]and Alexis Pokrovski [‡]

February 4, 2008

Abstract

Consider the operator $T = -\frac{d^2}{dx^2} + x^2 + q(x)$ in $L^2(\mathbb{R})$, where real functions q, q' and $\int_0^x q(s)\,ds$ are bounded. In particular, q is periodic or almost periodic. The spectrum of T is purely discrete and consists of the simple eigenvalues $\{\mu_n\}_{n=0}^{\infty}$, $\mu_n < \mu_{n+1}$. We determine their asymptotics $\mu_n = (2n+1) + (2\pi)^{-1}\int_{-\pi}^{\pi} q(\sqrt{2n+1}\sin\theta)\,d\theta + O(n^{-1/3})$.

1 Introduction and main results

Consider the quantum-mechanical harmonic oscillator $T^0 = -\frac{d^2}{dx^2} + x^2$ on $L^2(\mathbb{R})$. It is well known that the spectrum of T^0 is purely discrete and consists of the simple eigenvalues $\mu_n^0 = 2n+1$, $n \geqslant 0$ with corresponding orthonormed eigenfunctions ψ_n^0. Define the perturbed operator $Ty = -y'' + x^2 y + q(x)y$ in $L^2(\mathbb{R})$, where q belongs to the complex Banach space $\mathcal{B}$ given by

$$\mathcal{B} = \left\{ q, q', q_1 \in L^\infty(\mathbb{R}) : \|q\|_{\mathcal{B}} = \sup_{x\in\mathbb{R}} \Big(|q(x)| + |q'(x)| + |q_1(x)| \Big) < \infty \right\}, \quad q_1(x) \equiv \int_0^x q(t)dt. \tag{1.1}$$

In particular, this class includes periodic, almost-periodic q. If $q \in \mathcal{B}$ is real, then T is self-adjoint, its spectrum is purely discrete, $\sigma(T) = \{\mu_n\}_{n=0}^{\infty}$ and $\mu_n = \mu_n^0 + O(1),\ n \to \infty$. Our goal is to determine the asymptotics of $\mu_n - \mu_n^0$ as $n \to \infty$.

For decaying perturbations (e.g. $q', xq \in L^2(\mathbb{R})$) a complete inverse spectral theory is obtained in [2], [3]. At the same time we did not find in the literature any results concerning periodic and almost-periodic q. The existing methods (e.g. in [2]) cannot be used for $q \in \mathcal{B}$.

Let us number the points of the spectrum so that $|\mu_n| \leqslant |\mu_{n+1}|$ counting multiplicity.

---

[*] Institut für Mathematik, Universität Potsdam, Am Neuen Palais 10, 14469 Potsdam, Germany

[†] Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany

[‡] Institut für Mathematik, Universität Potsdam, and Laboratory of Quantum Networks, Institute for Physics, St.-Petersburg State University, Ulyanovskaya 1, 198504 St.-Petersburg, Russia.

Theorem 1.1. *For any $q \in \mathcal{B}$ the spectrum of operator T is discrete and the following asymptotic is fulfilled:*

$$\mu_n = \mu_n^0 + \mu_n^1 + \|q\|_{\mathcal{B}} O(n^{-\frac{1}{3}}), \qquad \mu_n^1 = \frac{1}{2\pi} \int_{-\pi}^{\pi} q(\sqrt{\mu_n^0} \sin \vartheta)\, d\vartheta = \|q\|_{\mathcal{B}} O(n^{-\frac{1}{4}}). \tag{1.2}$$

Remark. For finite smooth q the formula (1.2) gives $\mu_n^1 = \frac{1}{\pi\sqrt{\mu_n^0}} \int_{-\infty}^{\infty} q(s)\, ds$, which is the well-known leading term of asymptotics in this case [2].

Proposition 1.2. *Let $q \in \mathcal{B}$ and $q(x) = \int_{\mathbb{R}} e^{ixt}\, d\nu(t)$ for some Borel measure $d\nu$ on $\mathbb{R}$ which satisfies the condition $C_q = \int_{\mathbb{R}} (1 + |t|^{-p})\, d\nu(t) < \infty$ for some $p > \frac{3}{2}$. Then*

$$\mu_n^1 = \int_{\mathbb{R}} J_0(t\sqrt{\lambda})\, d\nu(t) = \frac{\sigma(\sqrt{\mu_n^0})}{(\mu_n^0)^{\frac{1}{4}}} + C_q O(n^{-\frac{3}{4}}), \tag{1.3}$$

where J_0 is the Bessel function and

$$\sigma(s) = \sqrt{\frac{2}{\pi}} \int_{\mathbb{R}} \frac{\cos(|t|s - \frac{\pi}{4})}{|t|^{\frac{1}{2}}} d\nu(t). \tag{1.4}$$

Moreover, if q has the form $q(x) = \sum_{k \in \mathbb{Z}} q_k e^{ixt_k}$, then

$$\sigma(s) = \sqrt{\frac{2}{\pi}} \sum_{k \in \mathbb{Z}} \frac{q_k}{\sqrt{|t_k|}} \cos(s|t_k| + \frac{\pi}{4}). \tag{1.5}$$

In Section 2 we introduce the quasiclassical change of variable. In this variable we write the integral equation for the fundamental solutions. In Sections 3 and 4 we prove convergence of the iteration series for the fundamental solutions in the sub-barrier ($x \gtrsim \sqrt{|\lambda|}$) and over-barrier ($0 < x \lesssim \sqrt{|\lambda|}$) regions, respectively. In Section 5 using these series we derive the asymptotics of the Wronskian. In Section 6 using this asymptotics we prove Theorem 1.1. We prove auxiliary properties of the quasiclassical change of variables in the Appendix.

2 Preliminaries: changes of variables

Consider the differential equation

$$-y'' + (x^2 + q(x))y = \lambda y, \qquad (x, \lambda) \in \mathbb{R} \times \mathbb{C}. \tag{2.1}$$

We shall show that there exist fundamental solutions $\psi_\pm$ which satisfy the asymptotics

$$\psi_\pm(x, \lambda) = (\pm\sqrt{2}x)^{\frac{\lambda-1}{2}} e^{-\frac{x^2}{2}} (1 + o(1)), \qquad \psi_\pm'(x, \lambda) = -x(\pm\sqrt{2}x)^{\frac{\lambda-1}{2}} e^{-\frac{x^2}{2}} (1 + o(1)) \tag{2.2}$$

as $x \to \pm\infty$ and locally uniformly in λ. If $q \equiv 0$ then these solutions have the form $\psi_\pm^0(x, \lambda) = D_{\frac{\lambda-1}{2}}(\pm\sqrt{2}x)$, where D_r is the Weber (parabolic cylinder) functions (see [1]). We introduce the Wronskian $\{f, g\} = fg' - f'g$.

Theorem 2.1. *Let $q, q_1 \in L^\infty(\mathbb{R})$. Then*
i) For any $\lambda \in \mathbb{C}$ there exist unique solutions $\psi_\pm(x,\lambda)$ of (2.1) with the asymptotics (2.2). Moreover, for each $x \in \mathbb{R}$ the functions $\psi_\pm(x,\cdot), \psi'_\pm(x,\cdot)$ and $w = \{\psi_-, \psi_+\}$ are entire.
ii) If q is real, then the operator $T = -\frac{d^2}{dx^2} + x^2 + q(x)$ has only simple eigenvalues.

Proof. Consider the function ψ_+, the proof for ψ_- is similar. In order to prove that ψ_+ is entire function of λ it is sufficient to show that it is analytic in each disc $\mathbb{D}(\mu) = \{\lambda \in \mathbb{C} : |\lambda - \mu| \leqslant 1\}, \mu \in \mathbb{C}$. For $\lambda \in \mathbb{D}(\mu)$ we have (see [1]) the uniform asymptotics

$$\psi_+^0(x,\lambda) = g(x)\left(1 + O\left(x^{-2}\right)\right), \quad {\psi_+^0}'(x,\lambda) = -xg(x)\left(1 + O\left(x^{-2}\right)\right), \quad x \to +\infty, \tag{2.3}$$

where $g(x) = (\sqrt{2}x)^{\frac{\lambda-1}{2}} e^{-\frac{x^2}{2}}$. Let $h(x,\lambda) = \frac{1}{2\sqrt{\pi}}\Gamma(\frac{1-\lambda}{2})(\psi_-^0(x,\lambda) - \sin\frac{\lambda}{2} \cdot \psi_+^0(x,\lambda))$; note that

$$h(x,\lambda) = \frac{1}{2xg(x)}(1 + O(x^{-1})), \quad h'(x,\lambda) = \frac{1}{2g(x)}(1 + O(x^{-1})), \quad x \to +\infty, \tag{2.4}$$

(see [1]) uniformly for $\lambda \in \mathbb{D}(\mu)$, so that $\{\psi_+^0, h\} = 1$. Define the entire function $M(x,y) = h(x,\lambda)\psi_0^+(y,\lambda) - \psi_0^+(x,\lambda)h(y,\lambda)$. Then a solution of

$$\psi(x,\lambda) = \psi_+^0(x,\lambda) + \lim_{t\to\infty}\int_x^t M(x,y)q(y)\psi(y,\lambda)\,dy \tag{2.5}$$

solves (2.1). We rewrite (2.5) in the form

$$p(x,\lambda) = p_0(x,\lambda) + \lim_{t\to\infty}\int_x^t K(x,y)q(y)p(y,\lambda)\,dy, \qquad x > 1, \tag{2.6}$$

where

$$p = \frac{\psi(x,\lambda)}{g(x)}, \quad p_0 = \frac{\psi_+^0(x,\lambda)}{g(x)}, \qquad K(x,y) = \frac{M(x,y)g(y)}{g(x)}. \tag{2.7}$$

Let $h_0 = 2xg(x)h(x,\lambda)$ and

$$K = U - V, \qquad U(x,y) = \frac{h_0(x)p_0(y)}{2x}\frac{g^2(y)}{g^2(x)}, \qquad V(x,y) = \frac{p_0(x)h_0(y)}{2y}. \tag{2.8}$$

In order to study Eq.(2.6) introduce the spaces of functions

$$\mathcal{F}_\alpha = \{f \in C([1,\infty)) : \|f\|_\alpha \equiv \sup_{x\in[1,\infty)} |x|^\alpha |f(x)| < \infty\}, \qquad \alpha \in \mathbb{R},$$

and $\mathcal{F}_{\alpha,\beta} = \{f \in \mathcal{F}_\alpha : f' \in \mathcal{F}_\beta\}$ with the norm $\|f\|_{\alpha,\beta} = \|f\|_\alpha + \|f'\|_\beta$, $\beta \in \mathbb{R}$. By (2.3), for $\lambda \in \mathbb{D}(\mu)$ we have the estimate

$$\|p_0\|_{0,1} \leqslant c < \infty. \tag{2.9}$$

Let $f \in \mathcal{F}_\alpha$ for some $\alpha \in \mathbb{R}$ and $u = Uqf$. Then we have

$$u(x) = \int_x^\infty \frac{h_0(x)p_0(y)}{2x}\frac{g^2(y)}{g^2(x)}q(y)f(y)dy = \frac{h_0(x)}{2x^\lambda}\int_x^\infty p_0(y)e^{x^2-y^2}y^{\lambda-1}q(y)f(y)dy. \tag{2.10}$$

Using (2.3), (2.4) and the estimate $\int_x^\infty e^{-y^2} y^\gamma dy \leqslant C e^{-x^2} x^{\gamma-1}$ for $x \geqslant 1$ and $\gamma \in \mathbb{R}$ we obtain

$$\|u\|_{\alpha+2} \leqslant C\|q\|_\infty \|f\|_\alpha, \qquad \|u'\|_{\alpha+1} \leqslant C\|q\|_\infty \|f\|_\alpha \tag{2.11}$$

uniformly in $\lambda \in \mathbb{D}(\mu)$. Here and below C is some absolute constant.

Let $f \in \mathcal{F}_{\alpha,\beta}$ for some $\alpha \in \mathbb{R}$ and $\beta > 0$. Set $v = Vqf$. Then integration by parts gives

$$v(x) = \lim_{t\to\infty} \int_x^t \frac{p_0(x)h_0(y)}{2y} q(y)f(y)\,dy = p_0(x)\int_x^\infty [q_1(x) - q_1(y)] \left(\frac{h_0(y)f(y)}{2y}\right)' dy,$$

where the last integral converges absolutely. Using (2.3–2.4) we obtain

$$|v(x)| \leqslant CC_q(\|f\|_\alpha x^{-\alpha-1} + \|f'\|_\beta x^{-\beta}), \qquad |v'(x)| \leqslant CC_q(\|f\|_\alpha x^{-\alpha-2} + \|f'\|_\beta x^{-\beta-1}) \tag{2.12}$$

for $x \geqslant 1$, uniformly in $\lambda \in \mathbb{D}(\mu)$, where $C_q = \|q\|_\infty + \|q_1\|_\infty$. Thus

$$K : \mathcal{F}_{\alpha,\beta} \to \mathcal{F}_{\alpha',\beta'}, \qquad \alpha' = \min\{\alpha+1, \beta\}, \quad \beta' = \min\{\alpha+1, \beta+1\}. \tag{2.13}$$

Consider the iterations $p_{n+1} = Kqp_n, n \geqslant 0$. By (2.9), we have $p_0 \in \mathcal{F}_{0,1}$; using (2.8), (2.11), (2.12) and (2.13), we conclude that

$$\|p_{n+1}\|_{\alpha_{n+1},\beta_{n+1}} \leqslant CC_q \|p_n\|_{\alpha_n,\beta_n}, \tag{2.14}$$

where $\alpha_0 = 0$, $\beta_0 = 1$,

$$\alpha_{2n} = \alpha_0 + n, \quad \beta_{2n} = \alpha_0 + 1 + n, \quad \alpha_{2n+1} = \alpha_0 + 1 + n, \quad \beta_{2n+1} = \alpha_0 + n. \tag{2.15}$$

Using (2.9) we obtain

$$|p_{2n}(x)| \leqslant (CC_q)^{2n} c x^{-n}, \quad |p_{2n}'(x)| \leqslant (CC_q)^{2n} c x^{-n-1},$$

$$|p_{2n+1}(x)| \leqslant (CC_q)^{2n+1} c x^{-n-1}, \quad |p_{2n+1}'(x)| \leqslant (CC_q)^{2n+1} c x^{-n-1}.$$

Hence for $x \geqslant x_0 = (2CC_q)^2$ the series $p(x) = \sum_{n\geqslant 0} p_n(x)$ and $p'(x) = \sum_{n\geqslant 0} p_n'(x)$ converge absolutely and uniformly in $\lambda \in \mathbb{D}(\mu)$; $p(x)$ gives the solution of Eq.(2.6). Moreover,

$$p(x) = 1 + O(x^{-1}), \qquad p'(x) = O(x^{-1}), \quad x \to \infty, \tag{2.16}$$

uniformly for $\lambda \in \mathbb{D}(\mu)$. Therefore $\psi = gp$ is a solution of (2.5). By (2.7) and (2.16), ψ satisfies (2.2). For each $n \geqslant 0$ and fixed $x \geqslant x_0$ the iterations $p_n(x,\cdot), p_n'(x,\cdot)$ are analytic in $\mathbb{D}(\mu)$. Hence $p(x,\cdot), p'(x,\cdot)$ are analytic in $\mathbb{D}(\mu)$ for each fixed $x \geqslant x_0$. By (2.7), $\psi_+(x,\lambda)$ and $\psi_+'(x,\lambda)$ are analytic in $\mathbb{D}(\mu)$ for each fixed $x \geqslant x_0$. Hence, the solution is also analytic in λ for any fixed x (see this simple fact e.g. in [4]). Thus $\psi_+(x,\lambda)$ and $\psi_+'(x,\lambda)$ are entire functions of λ for any $x \in \mathbb{R}$.

Suppose that $\psi_+(x,\lambda)$ is not unique and denote by $\widetilde{\psi}_+$ another solution of (2.1) satisfying (2.2). Then $\{\psi_+, \widetilde{\psi}_+\} = 0$ and therefore these solutions are linearly dependent. Since they have the same asymptotics (2.2), $\widetilde{\psi}_+ = \psi_+$. Thus ψ_+ is unique.

ii) Let λ be an eigenvalue, which is not simple. Then there exist solutions ψ_+ and ψ of (2.1) in $L^2(\mathbb{R})$. Note that ψ and ψ' are in $L^\infty(\mathbb{R})$. The asymptotics (2.2) yields $\{\psi_+, \psi\} = 0$. Therefore ψ_+ and ψ are linearly dependent, which gives contradiction. $\blacksquare$

Theorem 2.1 gives no information on high-energy asymptotics $\lambda \to \infty$. To derive those, we introduce some notations and auxiliary functions.

Throughout the paper we use the following agreements:

- *The functions* $\log z$ *and* $z^\alpha = e^{\alpha \log z}$ *for* $\alpha \in \mathbb{R}$ *take their principal values on* $\mathbb{C} \setminus \mathbb{R}_-$.
- *For* $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ *we set* $\lambda = |\lambda| e^{2i\vartheta}$, $\vartheta \in [0, \frac{\pi}{2}]$.

Define the function $\langle z \rangle = (1 + |z|^2)^{\frac{1}{2}}, z \in \mathbb{C}$. For any interval $I \subset \mathbb{R}$ we introduced a sector $S(I) = \{z \in \mathbb{C} : \arg z \in I\}$. Define the function

$$\xi(t) = \int_1^t \sqrt{s^2 - 1}\, ds = \frac{1}{2}\Big(t\sqrt{t^2 - 1} - \log(t + \sqrt{t^2 - 1})\Big), \quad t \in S(-\frac{\pi}{2}, 0), \tag{2.17}$$

where $\xi(t) > 0$ for $t > 1$. The function ξ is a conformal mapping from $S(-\frac{\pi}{2}, 0)$ onto $\Xi = \mathbb{C}_- \cup \{\operatorname{Re}\xi < 0, \operatorname{Im}\xi \in [0, \frac{\pi}{4})\}$. The following uniform asymptotics is fulfilled:

$$\xi(t) = \frac{1}{2}\Big(t^2 - \frac{1}{2} - \log 2t + O(|t|^{-2})\Big), \qquad |t| \to \infty, \qquad t \in S[-\frac{\pi}{2}, 0]. \tag{2.18}$$

We introduce the function

$$k(t) = \Big(\frac{3}{2}\xi(t)\Big)^{\frac{2}{3}}, \qquad t \in S(-\frac{\pi}{2}, 0), \quad k(0) = -\Big(\frac{3\pi}{8}\Big)^{\frac{2}{3}} < 0. \tag{2.19}$$

Note that $k(t)$ is a conformal mapping from $S(-\frac{\pi}{2}, 0)$ onto the domain $\mathcal{K}$ given by

$$\mathcal{K} = S[-\frac{2\pi}{3}, 0) \bigcup \Big\{k \in S(-\pi, -\frac{2\pi}{3}] : |k| \sin^{\frac{2}{3}} \frac{3 \arg k}{2} < |k(0)|\Big\}. \tag{2.20}$$

By (2.18), the following asymptotics and estimates are fulfilled:

$$k(t) = \Big(\frac{3}{4}\Big)^{\frac{2}{3}} t^{\frac{4}{3}}\Big(1 + O(t^{-1})\Big), \ \ t \in S[-\frac{\pi}{2}, 0], \qquad t(k) = \Big(\frac{4}{3}\Big)^{\frac{1}{2}} k^{\frac{3}{4}}\Big(1 + O(k^{-\frac{3}{4}})\Big), \ k \in \overline{\mathcal{K}}, \tag{2.21}$$

$$|t'(k)| \leqslant C\langle k \rangle^{-\frac{1}{4}}, \qquad |t''(k)| \leqslant C\langle k \rangle^{-\frac{5}{4}}, \quad k \in \overline{\mathcal{K}}, \tag{2.22}$$

where $t(k)$ is the inverse function for $k(t)$. Here and below C is an absolute constant.

Consider the change of variable $x \to k = k(\frac{x}{\sqrt{\lambda}})$; it maps $\mathbb{R}_+$ onto the curve $\tilde{\gamma}_\lambda = k(e^{i\vartheta}\mathbb{R}_+)$. The domain $\mathcal{K}$ and the curve $\tilde{\gamma}_\lambda$ are presented on Fig. 1. By (2.1), the function $y_1(k, \lambda) = \frac{y(\sqrt{\lambda} t(k))}{\sqrt{t'(k)}}$ solves

$$y_1''(k, \lambda) - \lambda^2 k y_1(k, \lambda) = v_0(k) y_1(k, \lambda) + v_q(k, \lambda) y_1(k, \lambda), \qquad k \in \tilde{\gamma}_\lambda, \tag{2.23}$$


$\mathcal{K}^* = k(S(0,\pi/2))$
$k-$plane
$k(0)$
$\mathcal{K} = k(S(-\pi/2,0))$
$\tilde\gamma_\lambda = k(e^{-i\vartheta}\mathbf{R}_+)$

Figure 1: The domain $\mathcal{K} = k(S(-\frac{\pi}{2},0))$ and the curve $\tilde\gamma_\lambda = k(e^{i\vartheta}\mathbb{R}_+)$.

where

$$v_0(k) = t'(k)^{\frac{3}{2}}\left(\frac{d^2}{dt^2}\frac{1}{\sqrt{k'(t)}}\right)\Big|_{t=t(k)}, \qquad v_q(k,\lambda) = \lambda\ t'(k)^2 q(\lambda^{\frac{1}{2}}t(k)). \tag{2.24}$$

Using (2.21) and (2.22) we obtain

$$|v_0(k)| \leqslant C(1+|k|)^{-2}, \qquad k \in \mathcal{K}. \tag{2.25}$$

For each $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ we define the **basic variable** $z = \lambda^{\frac{2}{3}}k$. We have the function

$$z(x,\lambda) = \lambda^{\frac{2}{3}}k(x\lambda^{-\frac{1}{2}}), \qquad \lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}, \qquad x \geqslant 0. \tag{2.26}$$

Each mapping $z(\cdot,\lambda) : \mathbb{R}_+ \to \Gamma_\lambda = z(\mathbb{R}_+,\lambda)$ is a real analytic isomorphism (see Fig. 2) and see Lemma 7.5 about Γ_λ. If $\lambda > 0$, then $\Gamma_\lambda = [-\lambda^{\frac{2}{3}}(\frac{3\pi}{8})^{\frac{2}{3}}, \infty)$ is a half-line. Moreover,

$$z_0 \equiv z(0,\lambda) = -\lambda^{\frac{2}{3}}\,(3\pi/8)^{\frac{2}{3}}, \qquad \lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}. \tag{2.27}$$

For any $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ and $0 \leqslant x_1 < x_2$ set $z_n = z(x_n,\lambda)$, $n = 1,2$. We define the curves

$$\Gamma_\lambda(z_1,z_2) = \{z : z = z(x,\lambda), x \in [x_1,x_2]\}, \quad \Gamma_\lambda(z_1) \equiv \Gamma_\lambda(z_1,\infty) = \{z : z = z(x,\lambda), x \geqslant x_1\}.$$

By (2.23), the function $u(z,\lambda) = y_1(z\lambda^{-\frac{2}{3}},\lambda)$ solves

$$\partial_z^2 u(z,\lambda) - zu(z,\lambda) = V(z,\lambda)u(z,\lambda), \quad V = V_0 + V_q, \qquad z \in \Gamma_\lambda, \tag{2.28}$$

here and below $\partial_z = \frac{\partial}{\partial z}$ and V_q (linear in q) and V_0 (not include q) are given by

$$V_0(z,\lambda) = \frac{v_0\left(z\lambda^{-\frac{2}{3}}\right)}{\lambda^{\frac{4}{3}}}, \quad V_q(z,\lambda) = \frac{\rho^2(z,\lambda)q\left(\sqrt{\lambda}t(z\lambda^{-\frac{2}{3}})\right)}{\lambda^{\frac{1}{3}}}, \quad \rho(z,\lambda) \equiv t'\left(z\lambda^{-\frac{2}{3}}\right). \tag{2.29}$$

Using (2.22), we obtain for $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ and $z \in S[-\pi, \frac{\pi}{3}]$ the estimates

$$|\rho(z,\lambda)| \leqslant C\langle z\lambda^{-\frac{2}{3}}\rangle^{-\frac{1}{4}}, \quad |\partial_z \rho(z,\lambda)| \leqslant |\lambda|^{-\frac{2}{3}}\langle z\lambda^{-\frac{2}{3}}\rangle^{-\frac{5}{4}}, \quad |V_0(z,\lambda)| \leqslant \frac{C}{|\lambda|^{\frac{4}{3}} + |z|^2}, \tag{2.30}$$

where C does not depend on λ and z.

To analyze (2.28) we need well-known properties [1] of the Airy functions Ai and Bi:

$$\mathrm{Ai}(z) = \frac{e^{-\frac{2}{3}z^{\frac{3}{2}}}}{2\sqrt[4]{z}}\Big(1 + O(z^{-\frac{3}{2}})\Big), \qquad |z| \to \infty, \qquad |\arg z| < \pi - \varepsilon, \quad \forall \varepsilon > 0, \tag{2.31}$$

$$\{\mathrm{Ai}(z), \mathrm{Bi}(z)\} = 1, \quad \mathrm{Bi}(z) = i\Big(2e^{-i\frac{\pi}{3}}\mathrm{Ai}(\omega z) - \mathrm{Ai}(z)\Big), \ \ \omega = e^{\frac{2\pi i}{3}}, \tag{2.32}$$

$$\mathrm{Ai}(z) = e^{-i\frac{\pi}{3}}\mathrm{Ai}(z\omega) + e^{i\frac{\pi}{3}}\mathrm{Ai}(z\overline{\omega}), \quad \mathrm{Bi}(z) = ie^{-i\frac{\pi}{3}}\mathrm{Ai}(z\omega) - ie^{i\frac{\pi}{3}}\mathrm{Ai}(z\overline{\omega}). \tag{2.33}$$

Let $\Gamma \subset \mathbb{C}$ be a smooth curve. For any continuous function f on Γ we denote by $\int_\Gamma f(s)\,ds$ the usual complex line integral. We denote by $\int_\Gamma f(s)\,|ds|$ the line integral of f along Γ with respect to the arc length $|ds| = \sqrt{(dx)^2 + (dy)^2}$. For integration along the infinite curve Γ_λ, defined above, we use the standard notation $\text{p.v.}\int_{\Gamma_\lambda(z)} f(s)\,ds = \lim \int_{\Gamma_\lambda(z,w)} f(s)\,ds$ as $w \to \infty, w \in \Gamma_\lambda^+$ whenever it exist.

We will study the formal integral equation

$$u_+(z,\lambda) = u_0(z) + \text{p.v.}\int_{\Gamma_\lambda(z)} J_0(z,s)V(s,\lambda)u_+(s,\lambda)\,ds, \quad z \in \Gamma_\lambda, \tag{2.34}$$

$$u_0(z) = \mathrm{Ai}(z), \qquad J_0(z,s) = \mathrm{Ai}(s)\mathrm{Bi}(z) - \mathrm{Ai}(z)\mathrm{Bi}(s), \quad z, s \in \mathbb{C}. \tag{2.35}$$

We rewrite (2.34) in the form

$$v_+(z) = a(z) + \text{p.v.}\int_{\Gamma_\lambda(z)} J(z,s)V(s,\lambda)v_+(s)\,ds, \quad a(z) \equiv \mathrm{Ai}(z)e^{\frac{2}{3}z^{\frac{3}{2}}}, \quad z \in \Gamma_\lambda, \tag{2.36}$$

$$u_+(z) = v_+(z)e^{-\frac{2}{3}z^{\frac{3}{2}}}, \qquad J(z,s) = J_0(z,s)e^{\frac{2}{3}(z^{\frac{3}{2}} - s^{\frac{3}{2}})}. \tag{2.37}$$

If $z < 0$ and $\lambda \in \overline{\mathbb{C}}_+$, then $z^{\frac{3}{2}}$ takes its values on the lower side of the cut. This agreement provides continuity as $\arg \lambda \downarrow 0$, since for $\lambda \in \mathbb{C}_+$ the curve Γ_λ lies in the lower half-plane. By (2.31) and (2.33), the following estimates are fulfilled:

$$|a(z)| \leqslant C\langle z\rangle^{-\frac{1}{4}}, \qquad \forall z \in \mathbb{C}, \tag{2.38}$$

$$|a'(z)| \leqslant C\langle z\rangle^{-\frac{5}{4}}, \quad |\arg z| \leqslant \pi - \varepsilon, \quad \forall \varepsilon > 0. \tag{2.39}$$

We write (2.36) in the form

$$v_+ = a + \mathbf{J}Vv_+, \tag{2.40}$$

where the integral operator $\mathbf{J}$ is given by

$$[\mathbf{J}f](z) = \text{p.v.} \int_{\Gamma_\lambda(z)} J(z,s)f(s)ds. \tag{2.41}$$

The next lemma (proved in the Appendix) gives a splitting of Γ_λ. Here and below we fix

$$\delta \in (0, \frac{4}{3} \arccos 2^{-\frac{1}{3}}). \tag{2.42}$$

Lemma 2.2. *For any $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ there exists a unique point $z_* \equiv z_*(\lambda) \in \Gamma_\lambda$ such that*

1. *if $0 \leqslant \arg \lambda < \delta$, then $|z_*| = \min_{z \in \Gamma_\lambda} |z|$,*
2. *if $\delta \leqslant \arg \lambda \leqslant \pi$, then $z_* = \Gamma_\lambda \cap \{z : \arg z = -\frac{\pi}{3}\}$.*

For any $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ here and below we use z_*, defined by Lemma 2.2. We define the point x_* by $z_* = z(x_*, \lambda)$ and let $t_* = \frac{x_*}{\sqrt{\lambda}}$ and set

$$\Gamma_\lambda^- = \Gamma_\lambda(z(0,\lambda), z_*), \qquad \Gamma_\lambda^+ = \Gamma_\lambda(z_*, \infty), \qquad \Gamma_\lambda = \Gamma_\lambda^- \cup \Gamma_\lambda^+. \tag{2.43}$$



Figure 2: The point z_* divides the curve $\Gamma_\lambda = \Gamma_\lambda^- \bigcup \Gamma_\lambda^+$ (for $\lambda = |\lambda| e^{2i\vartheta} \in \overline{\mathbb{C}}_+ \setminus \{0\}$).

Lemma 2.3. *Let $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$. Let $h(x,\lambda) = |\exp(\frac{2}{3} z(x,\lambda)^{\frac{3}{2}})|$ for $x \geqslant 0$. Then*

1. *if $\lambda > 0$, then $h(\cdot,\lambda)$ is strictly increasing on $[x_*, \infty)$ and $h(\cdot,\lambda) \equiv 1$ on $[0, x_*]$,*

2. if $0 < \arg\lambda \leqslant \pi$, *then* $h(\cdot,\lambda)$ *is strictly increasing on* $[0,\infty)$.

Lemma 2.4. *Let* $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ *and* $|\lambda| \geqslant 1$. *Assume a)* $z \in \Gamma_\lambda$, $\delta \leqslant \arg\lambda \leqslant \pi$ *or b)* $z \in \Gamma_\lambda^+$. *Then the following estimates are fulfilled:*

$$\int_{\Gamma_\lambda(z)} |e^{-\frac{4}{3}s^{\frac{3}{2}}}| \langle s\rangle^{-\alpha} |ds| \leqslant C |e^{-\frac{4}{3}z^{\frac{3}{2}}}| \langle z\rangle^{-\alpha-\frac{1}{2}}, \qquad \alpha \in \mathbb{R}, \tag{2.44}$$

$$\int_{\Gamma_\lambda(z)} \langle s\rangle^{-\alpha} |ds| \leqslant C \langle z\rangle^{-\alpha+1}, \qquad \alpha \geqslant 1, \tag{2.45}$$

where C *is independent of* λ *and* z.

Lemma 2.5. *Let* $\lambda \in \overline{\mathbb{C}}_+$ *and* $|\lambda| \geqslant 1$. *Then the following estimates are fulfilled:*

$$\int_{\Gamma_\lambda^-} \langle s\rangle^{-\alpha} |ds| \leqslant \begin{cases} C(1-\alpha)^{-1}|\lambda|^{\frac{2}{3}(1-\alpha)} & \text{for} \quad 0 \leqslant \alpha < 1, \\ C\log(|\lambda|+1) & \text{for} \quad \alpha = 1, \\ C(\alpha-1)^{-1} & \text{for} \quad \alpha > 1, \end{cases} \tag{2.46}$$

$$\int_{\Gamma_\lambda} \frac{|ds|}{|\lambda|^{\frac{4}{3}} + |s|^2} \leqslant \frac{C}{|\lambda|^{\frac{2}{3}}}, \tag{2.47}$$

where C *is independent of* λ *and* z.

3 Analysis of the integral equation

In this section we consider the integral equation $v_+ = a + \mathbf{J}Vv_+$ for large $|\lambda|$ in the following cases: i) $0 \leqslant \arg\lambda \leqslant \pi$, $z \in \Gamma_\lambda^+$ and ii) $\delta \leqslant \arg\lambda \leqslant \pi$, $z \in \Gamma_\lambda^-$. Moreover, we give the complete analysis for these cases. The case $0 \leqslant \arg\lambda \leqslant \pi$, $z \in \Gamma_\lambda^-$ is treated in the next section.

For $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ and $\alpha, \beta > 0$ define the Banach spaces of functions on Γ_λ:

$$\mathcal{F}_\alpha^\lambda = \Big\{ f \in C(\Gamma_\lambda^+) : \|f\|_\alpha \equiv \sup_{z\in\Gamma_\lambda^+} \langle z\rangle^\alpha |f(z)| < \infty \Big\} \qquad \text{for} \quad |\arg\lambda| < \delta, \tag{3.1}$$

$$\mathcal{F}_\alpha^\lambda = \Big\{ f \in C(\Gamma_\lambda) : \|f\|_\alpha \equiv \sup_{z\in\Gamma_\lambda} \langle z\rangle^\alpha |f(z)| < \infty \Big\} \qquad \text{for} \quad \delta \leqslant |\arg\lambda| \leqslant \pi, \tag{3.2}$$

$$\mathcal{F}_{\alpha,\beta}^\lambda = \Big\{ f \in F_\alpha^\lambda : f' \in F_\beta^\lambda \Big\}, \qquad \|f\|_{\alpha,\beta} = \|f\|_\alpha + \|f'\|_\beta. \tag{3.3}$$

Evidently $\mathcal{F}_\alpha^\lambda \subset \mathcal{F}_{\alpha'}^\lambda$ and $\mathcal{F}_{\alpha,\beta}^\lambda \subset \mathcal{F}_{\alpha',\beta'}^\lambda$ for $\alpha < \alpha'$ and $\beta < \beta'$. Now we formulate the main result of this section; its proof is given in the end of the section.

Theorem 3.1. *Let* $q, q_1 \in L^\infty(\mathbb{R})$, *and* $\lambda \in \overline{\mathbb{C}}_+$. *Then the equation* $v_+ = v_0 + \mathbf{J}Vv_+$, $v_0(z) \equiv a(z)$, *has a unique solution* $v_+ \in \mathcal{F}^\lambda_{\frac{1}{4},1}$ *for* $|\lambda|^{\frac{1}{6}} \geqslant 2c_0\left(\|q\|_\infty + \|q_1\|_\infty + 1\right)$, *where* $c_0 > 1$ *is an absolute constant. Moreover, the solution satisfies*

$$|v_+(z)| \leqslant C\langle z\rangle^{-\frac{1}{4}}, \qquad |v_+'(z)| \leqslant C\langle z\rangle^{-1}, \tag{3.4}$$

$$|v_+(z) - v_0(z)| \leqslant \varepsilon C\langle z\rangle^{-\frac{3}{4}}, \qquad |v_+'(z) - v_0'(z)| \leqslant \varepsilon C\langle z\rangle^{-1}, \tag{3.5}$$

where $\varepsilon = c_0|\lambda|^{-\frac{1}{6}}\left(\|q\|_\infty + \|q_1\|_\infty + 1\right)$.

We have the identity

$$\mathrm{Ai}(z\omega) = e^{\frac{2}{3}z^{\frac{3}{2}}}a(z\omega), \qquad \omega = e^{\frac{2\pi i}{3}}, \qquad -\pi \leqslant \arg z < \frac{\pi}{3}. \tag{3.6}$$

Using (2.32),(2.33) and (3.6) we write the kernel $J(z,s)$, given by (2.37), in terms of $a(z)$ and $a(\omega z)$. As a result we obtain

$$J(z,s) = -2ie^{-i\frac{\pi}{3}}\left(a(z)a(s\omega) - e^{\frac{4}{3}(z^{\frac{3}{2}} - s^{\frac{3}{2}})}a(z\omega)a(s)\right), \qquad z, s \in S(-\pi, \frac{\pi}{3}). \tag{3.7}$$

Note that by Lemma 7.5.1 , in both cases i) and ii) we have $\Gamma_\lambda(z) \subset S[-\pi + \frac{2}{3}\delta, 0]$, so (3.7) holds on $\Gamma_\lambda(z)$.

Following (3.7), we represent $\mathbf{J}V_q$ as the sum of two operators. In the next two Lemmas (3.2 and 3.3) we estimate these two operators in suitable functional spaces. In Lemma 3.4 we estimate $\mathbf{J}V_0$ (which is asymptotically small in comparison with $\mathbf{J}V_q$). These estimates, combined in Lemma 3.5, give an *a priori* estimate for $\mathbf{J}V$. In Theorem 3.1 we prove convergence of the iterations series for the equation $v_+ = v_0 + \mathbf{J}Vv_+$. This gives the estimates for v_+ necessary for further analysis.

For $v_0(z) \equiv a(z)$ given by (2.36), due to (2.38) and Lemma 7.5.1 we have

$$\|v_0\|_{\frac{1}{4},\frac{5}{4}} \leqslant C. \tag{3.8}$$

uniformly in $\lambda \in \overline{\mathbb{C}} \setminus \{0\}$. For any fixed $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$ and $z_1, z_2 \in \Gamma_\lambda$ such that $z_j = z(x_j, \lambda), j = 1, 2$ and $0 \leqslant x_1 \leqslant x_2$ we define the function

$$Q(z_2, z_1) \equiv \lambda^{-\frac{1}{6}} \int_{z_1}^{z_2} \hat{q}(z,\lambda)\rho(z,\lambda)\,dz, \tag{3.9}$$

where ρ is given by (2.29). We have the identity $Q(z_2, z_1) = \int_{x_2}^{x_1} q(x)\,dx$. Since $q_1(x) = \int_0^x q(t)dt \in L^\infty(\mathbb{R})$, we have

$$|Q(z_2, z_1)| \leqslant 2\|q_1\|_\infty \quad \text{for any} \quad z_1, z_2 \in \Gamma_\lambda. \tag{3.10}$$

By Lemma 7.5.2 and Lemma 7.5.3 , we have

$$C_1|\lambda|^{\frac{2}{3}} \leqslant |z| \leqslant C_2|\lambda|^{\frac{2}{3}} \quad \text{for} \quad z \in \Gamma_\lambda^-, \quad \delta \leqslant \arg\lambda \leqslant \pi, \tag{3.11}$$

where C_1 and C_2 are independent of z and λ.

Next we estimate the term of $\mathbf{J}V_q$, corresponding to the first term in decomposition (3.7).

Lemma 3.2. *Let $q, q_1 \in L^\infty(\mathbb{R})$. Assume $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$, $|\lambda| \geqslant 1$ and $f \in \mathcal{F}^\lambda_{\alpha,\beta}$ for $\alpha > 0, \beta > \frac{3}{4}$ (that is, f is defined on Γ_λ for $\delta \leqslant \arg\lambda \leqslant \pi$ and on Γ^+_λ for $0 \leqslant \arg\lambda \leqslant \delta$). Then $g(z,\lambda) = \mathrm{p.v.} \int_{\Gamma_\lambda(z)} a(s\omega)V_q(s)f(s)\,ds \in \mathcal{F}^\lambda_{\alpha,\beta}$ and satisfies*

$$|a(z)g(z,\lambda)| \leqslant C|\lambda|^{-\frac{1}{6}}\|q_1\|_\infty \Big\{ \|f\|_\alpha \langle z\rangle^{-\alpha-\frac{1}{2}} + \|f'\|_\beta \langle z\rangle^{-\beta+\frac{1}{2}} \Big\}, \tag{3.12}$$

$$\Big| a'(z)g(z,\lambda) \Big| \leqslant C|\lambda|^{-\frac{1}{6}}\|q_1\|_\infty \Big\{ \|f\|_\alpha \langle z\rangle^{-\alpha-\frac{3}{2}} + \|f'\|_\beta \langle z\rangle^{-\beta-\frac{1}{2}} \Big\}. \tag{3.13}$$

Proof. Consider the case $0 \leqslant \arg\lambda \leqslant \pi$, $z \in \Gamma^+_\lambda$. By Lemma 7.5.1 , we have $\Gamma_\lambda(z) \subset S[-\pi + \frac{2}{3}\delta, 0]$, so the uniform estimates (2.38) and (2.39) hold on $\Gamma_\lambda(z)$ for both $a(z)$ and $a(z\omega)$. Writing $F(z) = \lambda^{-\frac{1}{6}} a(z\omega)\rho(z,\lambda)$ (ρ is given by (2.29)), integration by parts yields

$$g(z,\lambda) = \mathrm{p.v.} \int_{\Gamma_\lambda(z)} (\partial_s Q(s,z))\, F(s)f(s)ds = -\mathrm{p.v.} \int_{\Gamma_\lambda(z)} Q(s,z)(F(s)f(s))'ds, \tag{3.14}$$

where we used $Q(z,z) = 0$ and $\lim\limits_{\Gamma^+_\lambda \ni w \to \infty} Q(w,z)F(w)f(w) = 0$ (this holds by (2.30), (2.38) and (3.10)). Thus using (2.30), (2.38), (2.39) and (3.10) we have

$$|g(z,\lambda)| \leqslant C\frac{\|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}} \Big\{ \int_{\Gamma_\lambda(z)} \|f\|_\alpha \langle s\rangle^{-\alpha-\frac{5}{4}} |ds|$$
$$+ \int_{\Gamma_\lambda(z)} \|f\|_\alpha |\lambda|^{-\frac{2}{3}} \langle \frac{s}{\lambda^{\frac{2}{3}}}\rangle^{-\frac{5}{4}} \langle s\rangle^{-\alpha-\frac{1}{4}} |ds| + \int_{\Gamma_\lambda(z)} \|f'\|_\beta \langle s\rangle^{-\beta-\frac{1}{4}} |ds| \Big\}.$$

Due to Lemma 7.5.4 we have $|z| = \inf\limits_{s\in\Gamma_\lambda(z)} |s|$. Using also (2.45) we obtain

$$|g(z,\lambda)| \leqslant C\frac{\|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}} \Big\{ \|f\|_\alpha \langle z\rangle^{-\alpha} \int_{\Gamma_\lambda(z)} \langle s\rangle^{-\frac{5}{4}} |ds| + \|f\|_\alpha \langle z\rangle^{-\alpha-\frac{1}{4}} \int_{\Gamma_\lambda(z)} |\lambda|^{-\frac{2}{3}} \langle \frac{s}{\lambda^{\frac{2}{3}}}\rangle^{-\frac{5}{4}} |ds|$$
$$+ \int_{\Gamma_\lambda(z)} \|f'\|_\beta \langle s\rangle^{-\beta-\frac{1}{4}} |ds| \Big\} \leqslant C|\lambda|^{-\frac{1}{6}}\|q_1\|_\infty \left\{ \|f\|_\alpha \langle z\rangle^{-\alpha-\frac{1}{4}} + \|f'\|_\beta \langle z\rangle^{-\beta+\frac{3}{4}} \right\}, \tag{3.15}$$

which together with (2.38) and (2.39) proves (3.12) and (3.13), respectively.

Consider the case $0 \leqslant \arg\lambda \leqslant \delta$, $z \in \Gamma^-_\lambda$. By Lemma 7.5.1 , we have $\Gamma_\lambda(z) \subset S[-\pi + \frac{2}{3}\delta, 0]$, so the uniform estimates (2.38) and (2.39) hold on $\Gamma_\lambda(z)$ for both $a(z)$ and $a(z\omega)$. We have $g(z,\lambda) = g_-(z,\lambda) + g_+(\lambda)$, where

$$g_+(\lambda) = \mathrm{p.v.} \int_{\Gamma_\lambda(z)} a(s\omega)V_q(s)f(s)\,ds, \quad g_-(z,\lambda) = \int_{\Gamma_\lambda(z,z_*)} a(s\omega)V_q(s)f(s)\,ds.$$

Using (3.11) and (3.15) for $z = z_*$ we have

$$|g_+(\lambda)| \leqslant C\frac{\|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}} \left\{ \frac{\|f\|_\alpha}{|z_*|^{\alpha+\frac{1}{4}}} + \frac{\|f'\|_\beta}{|z_*|^{\beta-\frac{3}{4}}} \right\} \leqslant C\frac{\|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}} \left\{ \frac{\|f\|_\alpha}{|z|^{\alpha+\frac{1}{4}}} + \frac{\|f'\|_\beta}{|z|^{\beta-\frac{3}{4}}} \right\}. \tag{3.16}$$

In order to estimate g_- we integrate by parts

$$g_-(z,\lambda) = Q(z_*,z)F(z)f(z) - \int_{\Gamma_\lambda(z,z_*)} Q(z_* s)\,(F(s)f(s))'\,ds, \quad F(z) = \lambda^{-\frac{1}{6}} a(z\omega)\rho(z,\lambda), \tag{3.17}$$

since $Q(z_*,z_*) = 0$ and $\hat{q}(s)\rho(s) = -\partial_s Q(z_*,s)$. Using (2.30), (2.38), (2.39) and (3.10) we obtain

$$|g_-(z,\lambda)| \leqslant C\frac{\|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}}\left\{\frac{\|f\|_\alpha}{|z|^{\alpha+\frac{1}{4}}} + \int_{\Gamma_\lambda^-}\left(\frac{\|f\|_\alpha}{|s|^{\alpha+\frac{5}{4}}} + \frac{\|f\|_\alpha}{|s|^{\alpha+\frac{1}{4}}|\lambda|^{\frac{2}{3}}}\right)|ds| + \int_{\Gamma_\lambda^-}\frac{\|f'\|_\beta}{|s|^{\beta+\frac{1}{4}}}|ds|\right\}.$$

By (2.46) and (3.11), we have

$$|g_-(z,\lambda)| \leqslant C\|q_1\|_\infty|\lambda|^{-\frac{1}{6}}\left\{\|f\|_\alpha|z|^{-\alpha-\frac{1}{4}} + \|f'\|_\beta|z|^{-\beta+\frac{3}{4}}\right\}. \tag{3.18}$$

Combining (3.16) and (3.18) with (2.38) gives (3.12). The estimate (3.13) follows from (3.16), (3.18) and (2.39). ■

For the analysis of the part of $\mathbf{J}V_q$ corresponding to the second term in (3.7) we also integrate by parts. Let us introduce an analogue of $Q(z_1,z_2)$:

$$P(z) = \frac{1}{\lambda^{\frac{1}{6}}}\int_{\Gamma_\lambda(z,\infty)} e^{-\frac{4}{3}s^{\frac{3}{2}}}\hat{q}(s,\lambda)\rho(s)ds = \int_x^\infty e^{-2\lambda\xi(s/\sqrt{\lambda})}q(s)\,ds, \quad z \in \Gamma_\lambda^+, \tag{3.19}$$

where $x = \sqrt{\lambda}\,t\left(\frac{z}{\lambda^{\frac{2}{3}}}\right)$. Using (2.30) and (2.44) for $q \in L^\infty(\mathbb{R})$ gives

$$|P(z)| \leqslant \|q\|_\infty C\int_{\Gamma_\lambda(z)}\frac{|e^{-\frac{4}{3}s^{\frac{3}{2}}}|}{(|\lambda|^{\frac{1}{6}} + |s|^{\frac{1}{4}})}|ds| \leqslant C\|q\|_\infty|e^{-\frac{4}{3}z^{\frac{3}{2}}}|\langle z\rangle^{-\frac{3}{4}}, \qquad z \in \Gamma_\lambda^+. \tag{3.20}$$

Using $|\rho(z)| \leqslant C$ and (2.44), we obtain another estimate

$$|P(z)| \leqslant \|q\|_\infty\frac{C}{|\lambda|^{\frac{1}{6}}}\int_{\Gamma_\lambda(z)}|e^{-\frac{4}{3}s^{\frac{3}{2}}}|\,|ds| \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{6}}}|e^{-\frac{4}{3}z^{\frac{3}{2}}}|\langle z\rangle^{-\frac{1}{2}}, \qquad z \in \Gamma_\lambda^+. \tag{3.21}$$

We estimate the part of $\mathbf{J}V_q$, corresponding to the second term in the decomposition (3.7).

Lemma 3.3. *Let $q \in L^\infty(\mathbb{R})$. Assume $\lambda \in \overline{\mathbb{C}}_+$, $|\lambda| \geqslant 1$, and $f \in \mathcal{F}_{\alpha,\beta}^\lambda$ for $\alpha > 0, \beta > 0$ (that is, f is defined on Γ_λ for $\delta \leqslant \arg\lambda \leqslant \pi$ and on Γ_λ^+ for $0 \leqslant \arg\lambda \leqslant \delta$). Then $g(z,\lambda) = \int_{\Gamma_\lambda(z)} a(s)e^{-\frac{4}{3}s^{\frac{3}{2}}}V_q(s)f(s)\,ds \in \mathcal{F}_{\alpha,\beta}^\lambda$ and satisfies*

$$|e^{\frac{4}{3}z^{\frac{3}{2}}}a(z)g(z,\lambda)| \leqslant C\|q\|_\infty|\lambda|^{-\frac{1}{6}}\left\{\|f\|_\alpha\langle z\rangle^{-\alpha-\frac{5}{4}} + \|f'\|_\beta\langle z\rangle^{-\beta-\frac{7}{4}}\right\}, \tag{3.22}$$

$$\left|g(z,\lambda)\frac{d}{dz}\Big(a(z\omega)e^{\frac{4}{3}z^{\frac{3}{2}}}\Big)\right| \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{6}}}\left\{\frac{\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{3}{4}}} + \frac{\|f'\|_\beta}{\langle z\rangle^{\beta+\frac{5}{4}}}\right\}. \tag{3.23}$$

Proof. Assume $0 \leqslant \arg\lambda \leqslant \pi$ and $z \in \Gamma_\lambda^+$. By Lemma 7.5.1 , we have $\Gamma_\lambda(z) \subset S[-\pi + \frac{2}{3}\delta, 0]$, so the uniform estimates (2.38) and (2.39) hold on $\Gamma_\lambda(z)$ for both $a(z)$ and $a(z\omega)$. Let $F(z) = \lambda^{-\frac{1}{6}} a(z)\rho(z)$, where ρ is given by (2.29).

$$g - PFf = I_1 + I_2, \quad I_1 = \int_{\Gamma_\lambda(z)} P(s)F'(s)f(s)ds, \qquad I_2 = \int_{\Gamma_\lambda(z)} P(s)F(s)f'(s)ds.$$

Using (2.30), (2.38) and (3.20) we have

$$|P(z)F(z)f(z)| \leqslant C|\lambda|^{-\frac{1}{6}}\|q\|_\infty |e^{-\frac{4}{3}z^{\frac{3}{2}}}| \|f\|_\alpha \langle z\rangle^{-\alpha-1}. \tag{3.24}$$

In order to estimate I_1 and I_2 we use (2.30), (2.38), (2.39), (2.44) and (3.21). This gives

$$|I_1| \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}} \int_{\Gamma_\lambda(z)} \left(\left\langle \frac{s}{\lambda^{\frac{2}{3}}} \right\rangle^{-\frac{1}{4}} \frac{\|f\|_\alpha |e^{-\frac{4}{3}s^{\frac{3}{2}}}|}{\langle s\rangle^{\alpha+\frac{7}{4}}} + \frac{|\lambda|^{-\frac{2}{3}}}{\left\langle \frac{s}{\lambda^{\frac{2}{3}}} \right\rangle^{\frac{5}{4}}} \frac{\|f\|_\alpha |e^{-\frac{4}{3}s^{\frac{3}{2}}}|}{\langle s\rangle^{\alpha+\frac{3}{4}}} \right) |ds|$$

$$\leqslant C\|q\|_\infty |\lambda|^{-\frac{1}{6}} |e^{-\frac{4}{3}z^{\frac{3}{2}}}| \|f\|_\alpha \langle z\rangle^{-\alpha-2},$$

$$|I_2| \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}} \int_{\Gamma_\lambda(z)} \|f'\|_\beta |e^{-\frac{4}{3}s^{\frac{3}{2}}}| \left\langle \frac{s}{\lambda^{\frac{2}{3}}} \right\rangle^{-\frac{1}{4}} \frac{|ds|}{\langle s\rangle^{\beta+\frac{3}{4}}} \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{6}}} |e^{-\frac{4}{3}z^{\frac{3}{2}}}| \frac{\|f'\|_\beta}{\langle z\rangle^{\beta+\frac{3}{2}}}.$$

The above estimates for I_1, I_2 and (3.24) give

$$|g(z,\lambda)| \leqslant C|\lambda|^{-\frac{1}{6}} \|q\|_\infty |e^{-\frac{4}{3}z^{\frac{3}{2}}}| \left\{ \|f\|_\alpha \langle z\rangle^{-\alpha-1} + \|f'\|_\beta \langle z\rangle^{-\beta-\frac{3}{2}} \right\}. \tag{3.25}$$

The last estimate together with (2.38) and (2.39) implies (3.22) and (3.23).

Assume $\delta \leqslant \arg\lambda \leqslant \pi$ and $z \in \Gamma_\lambda^-$. Using $\Gamma_\lambda = \Gamma_\lambda^- \cup \Gamma_\lambda^+$ we have $g = g_- + g_+$, where

$$g_-(z,\lambda) = \int_{\Gamma_\lambda(z,z_*)} a(s)e^{-\frac{4}{3}s^{\frac{3}{2}}} V_q(s)f(s)\,ds, \quad g_+(\lambda) = \int_{\Gamma_\lambda^+} a(s)e^{-\frac{4}{3}s^{\frac{3}{2}}} V_q(s)f(s)\,ds.$$

Using (3.11) and (3.25) for $z = z_*$ we obtain

$$|g_+(\lambda)| \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{6}}} |e^{-\frac{4}{3}z_*^{\frac{3}{2}}}| \left\{ \frac{\|f\|_\alpha}{|z_*|^{\alpha+1}} + \frac{\|f'\|_\beta}{|z_*|^{\beta+\frac{3}{2}}}, \right\} \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{6}}} |e^{-\frac{4}{3}z_*^{\frac{3}{2}}}| \left\{ \frac{\|f\|_\alpha}{|z|^{\alpha+1}} + \frac{\|f'\|_\beta}{|z|^{\beta+\frac{3}{2}}}, \right\}. \tag{3.26}$$

Using (2.38), (2.29), (2.30), (2.44) and (3.11) results in

$$|g_-(z,\lambda)| \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}} \int_{\Gamma_\lambda(z,z_*)} |e^{-\frac{4}{3}s^{\frac{3}{2}}}| \frac{\|f\|_\alpha}{|s|^{\alpha+\frac{1}{4}}} |ds| \leqslant C\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{6}}} |e^{-\frac{4}{3}z^{\frac{3}{2}}}| \frac{\|f\|_\alpha}{|z|^{\alpha+1}}. \tag{3.27}$$

Now (3.22) and (3.23) follow from (3.26) and (3.27) taking into account (2.38), (2.39) and the fact that, by Lemma 2.3, $|\exp\{\frac{2}{3}z(x,\lambda)^{\frac{3}{2}}\}|$ is strictly increasing. ■

In the following Lemma we estimate the operator $\mathbf{J}V_0$. We show that as $\lambda \to \infty$ it is asymptotically small in comparison with $\mathbf{J}V_q$.

Lemma 3.4. *Let $\lambda \in \overline{\mathbb{C}}_+$, $|\lambda| \geqslant 1$ and $f \in \mathcal{F}^\lambda_\alpha$ for some $\alpha > 0$ (that is, f is defined on Γ_λ for $\delta \leqslant \arg\lambda \leqslant \pi$ and on Γ^+_λ for $0 \leqslant \arg\lambda \leqslant \delta$). Then $\mathbf{J}V_0 f \in \mathcal{F}^\lambda_{\alpha+\frac{1}{2}}$ and*

$$|(\mathbf{J}V_0 f)(z,\lambda)| \leqslant \frac{C}{|\lambda|^{\frac{2}{3}}} \frac{\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{1}{2}}}, \qquad \left|\frac{\partial}{\partial z}(\mathbf{J}V_0 f)(z,\lambda)\right| \leqslant \frac{C}{|\lambda|^{\frac{2}{3}}} \frac{\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{3}{2}}}. \tag{3.28}$$

Proof. By Lemma 7.5.1 , we have $\Gamma_\lambda(z) \subset S[-\pi + \frac{2}{3}\delta, 0]$, so the decomposition (3.7) holds on $\Gamma_\lambda(z)$. We estimate the part of $\mathbf{J}V_0$, corresponding to the first term in decomposition (3.7). Using Lemma 7.5, (2.30), (2.38), (2.45) and the inequality $\langle z\rangle|\lambda|^{\frac{2}{3}} \leqslant 2(|\lambda|^{\frac{4}{3}} + |z|^2)$. As a result we have

$$\begin{aligned}\left|\int_{\Gamma_\lambda(z,\infty)} a(s\omega)V_0(s)f(s)\,ds\right| &\leqslant C\int_{\Gamma_\lambda(z,\infty)} \frac{\|f\|_\alpha}{\langle s\rangle^{\alpha+\frac{1}{4}}} \frac{|ds|}{|\lambda|^{\frac{4}{3}} + |s|^2} \\ &\leqslant \frac{C}{|\lambda|^{\frac{2}{3}}} \int_{\Gamma_\lambda(z,\infty)} \frac{\|f\|_\alpha}{\langle s\rangle^{\alpha+\frac{5}{4}}}|ds| \leqslant \frac{C}{|\lambda|^{\frac{2}{3}}} \frac{\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{1}{4}}}\end{aligned} \tag{3.29}$$

In order to estimate the part of $\mathbf{J}V_0$, corresponding to the second term in decomposition (3.7), we use (2.30), (2.38), (2.44) and the inequality $\langle z\rangle|\lambda|^{\frac{2}{3}} \leqslant 2(|\lambda|^{\frac{4}{3}} + |z|^2)$. This gives

$$\begin{aligned}\left|\int_{\Gamma_\lambda(z,\infty)} a(s)e^{-\frac{4}{3}s^{\frac{3}{2}}}V_0(s)f(s)\,ds\right| &\leqslant C\int_{\Gamma_\lambda(z,\infty)} \frac{\|f\|_\alpha}{|\lambda|^{\frac{4}{3}} + |s|^2} \frac{|e^{-\frac{4}{3}s^{\frac{3}{2}}}|}{\langle s\rangle^{\alpha+\frac{1}{4}}}\,|ds| \\ &\leqslant C\frac{\|f\|_\alpha}{|\lambda|^{\frac{2}{3}}} \int_{\Gamma_\lambda(z,\infty)} \frac{|e^{-\frac{4}{3}s^{\frac{3}{2}}}|}{\langle s\rangle^{\alpha+\frac{5}{4}}}\,|ds| \leqslant \frac{C}{|\lambda|^{\frac{2}{3}}} \frac{|e^{-\frac{4}{3}z^{\frac{3}{2}}}|\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{7}{4}}}.\end{aligned} \tag{3.30}$$

The first estimate in (3.28) follows from (3.29) and (3.30) taking into account (2.38) and (3.7).

In order to estimate $\partial_z g(z,\lambda)$ we note that $\partial_z g(z,\lambda) = \int_{\Gamma_\lambda(z)} \partial_z J(z,s)V_0(s)f(s)\,ds$. Therefore the second estimate in (3.28) follows from (3.29) and (3.30) taking into account (2.39) and (3.7). ■

Now in order to estimate the operator $\mathbf{J}V = \mathbf{J}V_q + \mathbf{J}V_0$ we combine the results of the three previous Lemmas.

Lemma 3.5. *Let $q, q_1 \in L^\infty(\mathbb{R})$. Assume $\lambda \in \overline{\mathbb{C}}_+$, $|\lambda| \geqslant 1$, and $f \in \mathcal{F}^\lambda_{\alpha,\beta}$ for $\alpha > 0, \beta > \frac{3}{4}$ (that is, f is defined on Γ_λ for $\delta \leqslant \arg\lambda \leqslant \pi$ and on Γ^+_λ for $0 \leqslant \arg\lambda \leqslant \delta$). Then $\mathbf{J}Vf \in \mathcal{F}^\lambda_{\alpha,\beta}$ and*

$$|(\mathbf{J}Vf)(z,\lambda)| \leqslant C\frac{\|q\|_\infty + \|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}} \left\{\frac{\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{1}{2}}} + \frac{\|f'\|_\beta}{\langle z\rangle^{\beta-\frac{1}{2}}}\right\} + \frac{C}{|\lambda|^{\frac{2}{3}}} \frac{\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{1}{2}}}, \tag{3.31}$$

$$|\partial_z(\mathbf{J}Vf)(z,\lambda)| \leqslant C\frac{\|q\|_\infty + \|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}} \left\{\frac{\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{3}{4}}} + \frac{\|f'\|_\beta}{\langle z\rangle^{\beta+\frac{1}{2}}}\right\} + \frac{C}{|\lambda|^{\frac{2}{3}}} \frac{\|f\|_\alpha}{\langle z\rangle^{\alpha+\frac{3}{2}}}. \tag{3.32}$$

Proof. Recall that $\mathbf{J}V = \mathbf{J}V_q + \mathbf{J}V_0$. By Lemma 7.5.1 , we have $\Gamma_\lambda(z) \subset S[-\pi + \frac{2}{3}\delta, 0]$, so the decomposition (3.7) holds on $\Gamma_\lambda(z)$. Taking into account this decomposition, we deduce that the combination of Lemmas 3.2 and 3.3 gives the estimate for $\mathbf{J}V_q$ (the corresponding terms in (3.31) and (3.32) contain curved brackets). Together with the estimate (3.28) for $\mathbf{J}V_0$ this proves (3.31) and (3.32). ■

Proof of Theorem 3.1. We present the proof for $\lambda \in \overline{\mathbb{C}}_+$, for $\lambda \in \overline{\mathbb{C}}_-$ it is analogous. Let $v_{n+1} = \mathbf{J}Vv_n, n \geqslant 0$. Substituting v_n, $g = v_{n+1}$ in (3.31), (3.32) and taking into account (3.8) for v_0 we obtain

$$|v_{n+1}(z)| \leqslant \frac{\varepsilon \|v_n\|_{\alpha_n}}{\langle z\rangle^{\alpha_n+\frac{1}{2}}} + \frac{\varepsilon \|v_n'\|_{\beta_n}}{\langle z\rangle^{\beta_n-\frac{1}{2}}}, \qquad |v_{n+1}'(z)| \leqslant \frac{\varepsilon \|v_n\|_{\alpha_n}}{\langle z\rangle^{\alpha_n+\frac{3}{4}}} + \frac{\varepsilon \|v_n'\|_{\beta_n}}{\langle z\rangle^{\beta_n+\frac{1}{2}}},$$

where

$$\alpha_0 = \frac{1}{4}, \beta_0 = \frac{5}{4}, \quad \alpha_{n+1} = \min\{\alpha_n + \frac{1}{2}, \beta_n - \frac{1}{2}\}, \quad \beta_{n+1} = \min\{\alpha_n + \frac{3}{4}, \beta_n + \frac{1}{2}\}. \tag{3.33}$$

Therefore

$$|v_{2n}(z)| \leqslant \varepsilon^{2n}\|v_0\|_{\frac{1}{4},\frac{5}{4}}\langle z\rangle^{-\frac{n}{4}-\frac{1}{4}}, \qquad |v_{2n+1}(z)| \leqslant \varepsilon^{2n+1}\|v_0\|_{\frac{1}{4},\frac{5}{4}}\langle z\rangle^{-\frac{n}{4}-\frac{3}{4}}, \tag{3.34}$$

$$|v_{2n}'(z)| \leqslant \varepsilon^{2n}\|v_0\|_{\frac{1}{4},\frac{5}{4}}\langle z\rangle^{-\frac{n}{4}-\frac{5}{4}}, \quad |v_{2n+1}'(z)| \leqslant \varepsilon^{2n+1}\|v_0\|_{\frac{1}{4},\frac{5}{4}}\langle z\rangle^{-\frac{n}{4}-1}, \tag{3.35}$$

and for $\varepsilon < 1$ the series $v_+(z) = \sum_{n=0}^\infty v_n(z)$ converges absolutely and is a solution of $v_+ = a + \mathbf{J}Vv_+$. For $\varepsilon < \frac{1}{2}$ we obtain from (3.8), (3.34) and (3.35) the estimates (3.4), (3.5).

We prove the uniqueness. Suppose that there exists another solution $v_+^{(1)} \in \mathcal{F}^\lambda_{\frac{1}{4},1}$ and let $y = v_+ - v_+^{(1)} \in \mathcal{F}^\lambda_{\frac{1}{4},1}$. We have $y = \mathbf{J}Vy$ and therefore $y = (\mathbf{J}V)^n y$ for any integer $n \geqslant 1$; applying Lemma 3.5 we obtain $|y(z)| \leqslant C\varepsilon^n \|y\|_{\frac{1}{4},1}$. Taking the limit $n \to \infty$ for $\varepsilon < 1$ we obtain $y = 0$. ■

4 Uniform asymptotics

In this section we consider the equation $v_+ = a + \mathbf{J}v_+$ for large $|\lambda|$, $|\arg\lambda| \leqslant \delta$ and $z \in \Gamma_\lambda^-$. The case $z \in \Gamma_\lambda^+$ was treated in the previous section.

For $|\arg\lambda| \leqslant \delta$ denote by $\mathcal{F}^\lambda_-$ the class of functions f on Γ_λ^- such that $f, f' \in L^\infty(\Gamma_\lambda^-)$. We also set $\mathcal{F}^\lambda = \mathcal{F}^\lambda_- \oplus \mathcal{F}^\lambda_{\frac{1}{4},1}$ (for the definition of $\mathcal{F}^\lambda_{\alpha,\beta}$ see (3.3)). Our main result is

Theorem 4.1. *Let $q \in \mathcal{B}$ and $|\arg\lambda| \leqslant \delta$. Then the equation $v_+ = v_0 + \mathbf{J}Vv_+$, $v_0 \equiv a$, has a unique solution $v_+ \in \mathcal{F}^\lambda$ for $|\lambda|^{\frac{1}{6}} \geqslant 2c_0(\|q\|_{\mathcal{B}} + 1)$, where $c_0 > 1$ is an absolute constant. If, in addition, $(z, \lambda) \in \Gamma_\lambda^- \times S[-\delta, \delta]$, then the following estimates are fulfilled:*

$$|v_+(z)| \leqslant C\langle z\rangle^{-\frac{1}{4}}, \qquad |v_+'(z)| \leqslant C\langle z\rangle^{\frac{1}{4}}, \tag{4.1}$$

$$|v_1(z)|\leqslant C\varepsilon\langle z\rangle^{-\frac{1}{4}}, \qquad |v_1'(z)|\leqslant C\langle z\rangle^{\frac{1}{4}}\varepsilon, \qquad v_1=\mathbf{J}Vv_0, \qquad \varepsilon=c_0|\lambda|^{-\frac{1}{6}}(\|q\|_{\mathcal{B}}+1), \tag{4.2}$$

$$|v_+(z)-v_0(z)-v_1(z)|\leqslant C\varepsilon^2\langle z\rangle^{-\frac{1}{4}}, \qquad |v_+'(z)-v_0'(z)-v_1'(z)|\leqslant \varepsilon^2 C\langle z\rangle^{\frac{1}{4}}. \tag{4.3}$$

Corollary 4.2. *Let* $q\in\mathcal{B}$. *Then the equation* $u_+=\mathrm{Ai}+\mathbf{J_0}Vu_+$ *has a unique solution* $u_+(z,\lambda)$ *such that* $u_+(z,\lambda)=\mathrm{Ai}(z)(1+o(1))$, $\partial_z u_+(z,\lambda)+\sqrt{z}u_+(z,\lambda)=\mathrm{Ai}'(z)O(z^{-\frac{5}{4}})$ *as* $\Gamma_\lambda\ni z\to\infty$ *for* $|\lambda|^{\frac{1}{6}}\geqslant 2c_0(\|q\|_{\mathcal{B}}+1)$, *where* $c_0>1$ *is an absolute constant. Moreover,* $u_+(z,\lambda)=e^{-\frac{2}{3}z^{\frac{3}{2}}}v_+(z,\lambda)$. *The following estimates for* u_+ *and* $u_1(z,\lambda)=e^{-\frac{2}{3}z^{\frac{3}{2}}}v_1(z,\lambda)$ *are fulfilled uniformly in* $z\in\Gamma_\lambda^-$*:*
If $|\arg\lambda|\leqslant\delta$, *then*

$$|u_+(z,\lambda)|\leqslant Ce^{\frac{2}{3}|\operatorname{Re}z^{\frac{3}{2}}|}\langle z\rangle^{-\frac{1}{4}}, \qquad |\partial_z u_+(z,\lambda)|\leqslant C\langle z\rangle^{\frac{1}{4}}e^{\frac{2}{3}|\operatorname{Re}z^{\frac{3}{2}}|}, \tag{4.4}$$

$$|u_1(z,\lambda)|\leqslant C\varepsilon\frac{e^{\frac{2}{3}|\operatorname{Re}z^{\frac{3}{2}}|}}{\langle z\rangle^{\frac{1}{4}}}, \quad |\partial_z u_1(z,\lambda)|\leqslant C\varepsilon\langle z\rangle^{\frac{1}{4}}e^{\frac{2}{3}|\operatorname{Re}z^{\frac{3}{2}}|}, \quad \varepsilon=c_0\frac{\|q\|_{\mathcal{B}}+1}{|\lambda|^{\frac{1}{6}}}, \tag{4.5}$$

$$|u_+(z,\lambda)-\mathrm{Ai}(z)-u_1(z,\lambda)|\leqslant C\varepsilon^2e^{\frac{2}{3}|\operatorname{Re}z^{\frac{3}{2}}|}\langle z\rangle^{-\frac{1}{4}}, \tag{4.6}$$

$$|\partial_z u_+(z,\lambda)-\mathrm{Ai}'(z)-\partial_z u_1(z,\lambda)|\leqslant C\varepsilon^2\langle z\rangle^{\frac{1}{4}}e^{\frac{2}{3}|\operatorname{Re}z^{\frac{3}{2}}|}, \tag{4.7}$$

if $\delta\leqslant|\arg\lambda|\leqslant\pi$, *then*

$$|u_+(z,\lambda)-\mathrm{Ai}(z)|\leqslant C\varepsilon e^{\frac{2}{3}|\operatorname{Re}z^{\frac{3}{2}}|}\langle z\rangle^{-\frac{3}{4}}, \qquad |\partial_z u_+(z,\lambda)-\mathrm{Ai}'(z)|\leqslant C\varepsilon e^{\frac{2}{3}|\operatorname{Re}z^{\frac{3}{2}}|}\langle z\rangle^{-\frac{1}{4}}. \tag{4.8}$$

Proof. Set $u_+(z,\lambda)=e^{-\frac{2}{3}z^{\frac{3}{2}}}v_+(z,\lambda)$, where v_+ is given by Theorem 3.1 (for $\delta<|\arg\lambda|\leqslant\pi$) and Theorem 4.1 (for $|\arg\lambda|\leqslant\delta$). By (2.36), (2.37) and (3.4), u_+ is a solution of (2.34) with the required asymptotics.

In order to prove uniqueness, we suppose that there exists another solution $u_+^{(1)}$ with the same asymptotics as $\Gamma_\lambda\ni z\to\infty$. Then $v_+^{(1)}(z,\lambda)=e^{\frac{2}{3}z^{\frac{3}{2}}}u_+^{(1)}(z,\lambda)$ is in $\mathcal{F}^\lambda$ (for $|\arg\lambda|\leqslant\delta$) or in $\mathcal{F}^\lambda_{\frac{1}{4},1}$ (for $\delta<|\arg\lambda|\leqslant\pi$) and solves $v_+=v_0+\mathbf{J}Vv_+$. Hence, by Theorems 3.1 and 4.1, $v_+^{(1)}=v_+$, implying $u_+^{(1)}=u_+$.

The estimates (4.4–4.7) follow from (4.1–4.3). The estimate (4.8) follows from (3.5). ■

Below we consider only the case $0\leqslant\arg\lambda\leqslant\delta$, the case $-\delta\leqslant\arg\lambda\leqslant 0$ is analogous.

By Lemma 7.5.1 , Γ_λ^- is arbitrarily close to $\mathbb{R}_-$ as $\arg\lambda\to 0$. Thus we cannot use (2.39) in order to estimate the terms in the decomposition (3.7) for $J(z,s)$. So we use another representation. We have

$$\mathrm{Ai}(z\overline{\omega})=e^{-\frac{2}{3}z^{\frac{3}{2}}}a(z\overline{\omega}), \qquad z\in S[-\pi,-\frac{\pi}{3}), \qquad \omega=e^{\frac{2\pi i}{3}}. \tag{4.9}$$

Using (2.32), (2.33) and (3.6), we obtain from (2.35) and (2.37)

$$J(z,s) = -2i\left(a(z\overline{\omega})a(s\omega) - e^{\frac{4}{3}z^{\frac{3}{2}}-\frac{2}{3}s^{\frac{3}{2}}}a(z\omega)\mathrm{Ai}(s\overline{\omega})\right), \qquad z \in \Gamma_\lambda^-, \quad s \in \Gamma_\lambda, \tag{4.10}$$

where, by Lemma 7.3.1 , $\Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{2} + \frac{5}{12}\delta] \subset S[-\pi, -\frac{\pi}{3})$ and $\Gamma_\lambda^+ \subset S[-\frac{\pi}{2} - \frac{\delta}{4}, 0] \subset S(-\pi, \pi]$.

As it follows from the decomposition (4.10), the formal operator $\mathbf{J}V$ can be presented as the sum $a(z\overline{\omega})\times$(integral operator)$+e^{\frac{4}{3}z^{\frac{3}{2}}}a(z\omega)\times$(integral operator). Thus in order to estimate $\mathbf{J}V$ we estimate these integral operators, introducing for functions on Γ_λ^- a decomposition in the spirit of (4.10).

For $f \in \mathcal{F}_-^\lambda$ and a fixed decomposition

$$f(z) = a(z\overline{\omega})f_p(z) + e^{\frac{4}{3}z^{\frac{3}{2}}}a(z\omega)f_e(z), \quad z \in \Gamma_\lambda^-, \quad \lambda \in S[0,\delta]\bigcap\{\lambda \in \mathbb{C} : |\lambda| \geqslant 1\}, \tag{4.11}$$

we introduce the functionals

$$p_0(f,\lambda) = \sup_{z\in\Gamma_\lambda^-}\left(|f_p(z)| + |e^{\frac{4}{3}z^{\frac{3}{2}}}f_e(z)|\right), \quad p_1(f,\lambda) = \sup_{z\in\Gamma_\lambda^-}\langle z\rangle^{\frac{1}{2}}\left(|f_p'(z)| + |e^{\frac{4}{3}z^{\frac{3}{2}}}f_e'(z)|\right). \tag{4.12}$$

If $f \in \mathcal{F}_-^\lambda$ has a decomposition (4.11), then we have

$$|f(z)| \leqslant Cp_0(f,\lambda)\langle z\rangle^{-\frac{1}{4}}, \qquad |f'(z)|\langle z\rangle^{-\frac{1}{4}} \leqslant C\left(p_0(f,\lambda) + p_1(f,\lambda)\langle z\rangle^{-1}\right), \quad z \in \Gamma_\lambda^-. \tag{4.13}$$

If $f \in \mathcal{F}_-^\lambda$ has a decomposition (4.11), then we have

$$|f_p(z)| \leqslant p_0(f,\lambda), \qquad |f_e(z)| \leqslant |e^{-\frac{4}{3}z^{\frac{3}{2}}}|p_0(f,\lambda), \tag{4.14}$$

$$|f_p'(z)| \leqslant p_1(f,\lambda)\langle z\rangle^{-\frac{1}{2}}, \qquad |f_e'(z)| \leqslant |e^{-\frac{4}{3}z^{\frac{3}{2}}}|\langle z\rangle^{-\frac{1}{2}}p_1(f,\lambda). \tag{4.15}$$

Using (3.6), (4.9) and the identity (2.33) we obtain for $0 \leqslant \arg\lambda \leqslant \delta$

$$v_0(z) = a(z) = e^{i\frac{\pi}{3}}a(z\overline{\omega}) + e^{-i\frac{\pi}{3}}e^{\frac{4}{3}z^{\frac{3}{2}}}a(z\omega), \qquad z \in \Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3}). \tag{4.16}$$

For this decomposition using (2.38) and (2.39) we obtain

$$p_0(v_0,\lambda) \leqslant 2, \qquad p_1(v_0,\lambda) = 0, \qquad 0 \leqslant \arg\lambda \leqslant \delta. \tag{4.17}$$

Below we estimate $(\mathbf{J}Vf)\Big|_{\Gamma_\lambda^-}$ in terms of p_0 and p_1 assuming that $f \in \mathcal{F}^\lambda = \mathcal{F}_-^\lambda \oplus \mathcal{F}_{\frac{1}{4},1}^\lambda$. Using (4.10) for some function f on Γ_λ^- we obtain the following formal decomposition:

$$(\mathbf{J}Vf)(z) = a(z\overline{\omega})g_p(z) + e^{\frac{4}{3}z^{\frac{3}{2}}}a(z\omega)g_e(z), \quad g_p = \mathbf{J}_pVf, \quad g_e = \mathbf{J}_eVf, \quad z \in \Gamma_\lambda^-, \tag{4.18}$$

where $\lambda \in S[0,\delta] \bigcap \{\lambda \in \mathbb{C} : |\lambda| \geqslant 1\}$ and the operators $\mathbf{J}_p$ and $\mathbf{J}_e$ are given by

$$(\mathbf{J}_p f)(z) = -2i \int_{\Gamma_\lambda(z)} a(s\omega) f(s)\, ds, \quad (\mathbf{J}_e f)(z) = 2i \int_{\Gamma_\lambda(z)} e^{-\frac{2}{3}s^{\frac{3}{2}}} \mathrm{Ai}(s\overline{\omega}) f(s)\, ds. \tag{4.19}$$

In the applications below the integral along the infinite curve Γ_λ^+ exists in the principal value sense. In this section we will always define $p_0(\mathbf{J}Vf,\lambda)$ and $p_1(\mathbf{J}Vf,\lambda)$ using the decomposition (4.18) and (4.19).

In order to estimate $\mathbf{J}_pV$ and $\mathbf{J}_eV$ we decompose the integrals in (4.19) corresponding to the splitting $\Gamma_\lambda(z) = \Gamma_\lambda(z,z_*) \cup \Gamma_\lambda^+$. Thus we have

$$(\mathbf{J}_p f)(z) = (\mathbf{j}_p f)(z) + h_p(f), \qquad z \in \Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3}), \tag{4.20}$$

where, by definition

$$(\mathbf{j}_p f)(z) = -2i \int_{\Gamma_\lambda(z,z_*)} a(s\omega) f(s)\, ds, \quad h_p(f) = -2i \int_{\Gamma_\lambda^+} a(s\omega) f(s)\, ds. \tag{4.21}$$

A similar decomposition of $\mathbf{J}_e$ gives

$$(\mathbf{J}_e f)(z) = (\mathbf{j}_e f)(z) + h_e(f), \qquad z \in \Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3}), \tag{4.22}$$

where, by definition

$$(\mathbf{j}_e f)(z) = 2i \int_{\Gamma_\lambda(z,z_*)} e^{-\frac{2}{3}s^{\frac{3}{2}}} \mathrm{Ai}(s\overline{\omega}) f(s)\, ds, \quad h_e(f) = 2i \int_{\Gamma_\lambda^+} e^{-\frac{2}{3}s^{\frac{3}{2}}} \mathrm{Ai}(s\overline{\omega}) f(s)\, ds. \tag{4.23}$$

Note that the standard asymptotics (2.31) for $\mathrm{Ai}(s\overline{\omega})$ fails in the neighborhood of $\arg s = -\frac{\pi}{3}$. Taking into account (4.9) we obtain

$$(\mathbf{j}_e f)(z) = 2i \int_{\Gamma_\lambda(z,z_*)} e^{-\frac{4}{3}s^{\frac{3}{2}}} a(s\overline{\omega}) f(s)\, ds. \tag{4.24}$$

By (2.33), (2.36), (3.6) and (4.9), for $s \in \Gamma_\lambda^+ \subset S(-\pi, \frac{\pi}{3})$ we have $e^{-\frac{2}{3}s^{\frac{3}{2}}} \mathrm{Ai}(s\overline{\omega}) = e^{i\frac{\pi}{3}} a(s\omega) - e^{-\frac{4}{3}s^{\frac{3}{2}}} \omega a(s)$. This gives

$$h_e(f) = 2i \int_{\Gamma_\lambda^+} \left(e^{i\frac{\pi}{3}} a(s\omega) - e^{-\frac{4}{3}s^{\frac{3}{2}}} \omega a(s) \right) f(s)\, ds. \tag{4.25}$$

Using (4.19–4.25), for $z \in \Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3})$ we have the decomposition

$$(\mathbf{J}f)(z) = a(z\overline{\omega}) \left((\mathbf{j}_p f)(z) + h_p(f) \right) + e^{\frac{4}{3}z^{\frac{3}{2}}} a(z\omega) ((\mathbf{j}_e f)(z) + h_e(f)). \tag{4.26}$$

For $z_j = z(x_j, \lambda) \in \Gamma_\lambda^-$, $j = 1,2$ and $\quad x_1 < x_2$ we define

$$P_\pm(z_1, z_2) = \lambda^{\frac{1}{6}} \int_{\Gamma_\lambda(z_1,z_2)} e^{\pm\frac{4}{3}s^{\frac{3}{2}}} \rho^{-1}(s) V_q(s)\, ds. \tag{4.27}$$

Lemma 4.3. *Let* $q \in \mathcal{B}$. *Assume* $0 \leqslant \arg\lambda \leqslant \delta$, $|\lambda| \geqslant 1$. *Assume that* $f\big|_{\Gamma_\lambda^-} \in \mathcal{F}_-^\lambda$ *has a decomposition (4.11) and* $f\big|_{\Gamma_\lambda^+} = 0$. *Then* $(\mathbf{J}V_q f)(z) = a(z\overline{\omega})(\mathbf{j}_p V_q f)(z) + e^{\frac{4}{3}z^{\frac{3}{2}}} a(z\omega)(\mathbf{j}_e V_q f)(z) \in \mathcal{F}_-^\lambda$ *and for this decomposition the following estimates are fulfilled:*

$$p_0(\mathbf{J}V_q f, \lambda) \leqslant C\|q\|_{\mathcal{B}}|\lambda|^{-\frac{1}{6}}\Big(p_0(f,\lambda) + p_1(f,\lambda)\log(|\lambda|+1)\Big), \tag{4.28}$$

$$p_1(\mathbf{J}V_q f, \lambda) \leqslant C\|q\|_\infty|\lambda|^{-\frac{1}{3}} p_0(f,\lambda). \tag{4.29}$$

Proof. By Lemma 7.3.1 , $\Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3})$, so (4.10) holds. Using (4.11), (4.18), (4.20), (4.21), (4.22), (4.23) and (4.24) we obtain

$$(\mathbf{j}_p V_q f)(z) = -2i(I_{pp}(z) + I_{pe}(z)), \qquad (\mathbf{j}_e V_q f)(z) = 2i(I_{ep}(z) + I_{ee}(z)), \tag{4.30}$$

where

$$I_{pp}(z) = \int_{\Gamma_\lambda(z,z_*)} a(s\omega)a(s\overline{\omega})V_q(s)f_p(s)\,ds, \quad I_{pe}(z) = \int_{\Gamma_\lambda(z,z_*)} a(s\omega)^2 e^{\frac{4}{3}s^{\frac{3}{2}}} V_q(s)f_e(s)\,ds,$$

$$I_{ep}(z) = \int_{\Gamma_\lambda(z,z_*)} a(s\overline{\omega})^2 e^{-\frac{4}{3}s^{\frac{3}{2}}} V_q(s)f_p(s)\,ds, \quad I_{ee}(z) = \int_{\Gamma_\lambda(z,z_*)} a(s\omega)a(s\overline{\omega})V_q(s)f_e(s)\,ds.$$

We estimate I_{pp}. Define the functions $F_j(s) = \lambda^{-\frac{1}{6}} a(s\omega)a(s\overline{\omega})f_j(s)\rho(s)$ for $j = e, p$. Integration by parts yields

$$I_{pp}(z) = -Q(z_*, z)F_p(z) + \int_{\Gamma_\lambda(z,z_*)} Q(z_*, s)F_p'(s)ds, \tag{4.31}$$

where Q is given by (3.9). Using (2.29), (2.30), (2.38), (2.39) and (3.10), we obtain

$$|I_{pp}(z)| \leqslant C\frac{\|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}}\left[\frac{|f_p(z)|}{\langle z\rangle^{\frac{1}{2}}} + \int_{\Gamma_\lambda^-}\left\{\frac{|f_p'(s)|}{\langle s\rangle^{\frac{1}{2}}} + \left(\frac{|f_p(s)|}{\langle s\rangle^{\frac{3}{2}}} + \frac{|\lambda|^{-\frac{2}{3}}|f_p(s)|}{\langle s\rangle^{\frac{1}{2}}}\right)\right\}|ds|\right].$$

Due to (2.46), (4.14) and (4.15) we have

$$|I_{pp}(z)| \leqslant C\|q_1\|_\infty|\lambda|^{-\frac{1}{6}}\Big(p_0(f,\lambda) + p_1(f,\lambda)\log(|\lambda|+1)\Big), \quad z \in \Gamma_\lambda^-. \tag{4.32}$$

We estimate $I_{pp}'(z) = -a(z\omega)a(z\overline{\omega})V_q(z)f_p(z)$. Using (2.29), (2.30) and (2.38) we have

$$\big|I_{pp}'(z)\big| \leqslant C\|q\|_\infty|f_p(z)||\lambda|^{-\frac{1}{3}}\langle z\rangle^{-\frac{1}{2}}, \quad z \in \Gamma_\lambda^-. \tag{4.33}$$

The estimate of I_{ee} is similar. Using F_e and Q we integrate by parts. Using (2.29), (2.30), (2.38), (2.39) and (3.10), we obtain

$$|I_{ee}(z)| \leqslant C\frac{\|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}}\left[\frac{|f_e(z)|}{\langle z\rangle^{\frac{1}{2}}} + \int_{\Gamma_\lambda^-}\left\{\frac{|f_e'(s)|}{\langle s\rangle^{\frac{1}{2}}} + \left(\frac{|f_e(s)|}{\langle s\rangle^{\frac{3}{2}}} + \frac{|\lambda|^{-\frac{2}{3}}|f_e(s)|}{\langle s\rangle^{\frac{1}{2}}}\right)\right\}|ds|\right]$$

We have $|e^{-\frac{4}{3}z^{\frac{3}{2}}}| \geqslant 1$ for $z \in \Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3})$. Using (2.46), (4.14) and (4.15) we have

$$|I_{ee}(z)| \leqslant C\frac{\|q_1\|_\infty}{|\lambda|^{\frac{1}{6}}}|e^{-\frac{4}{3}z^{\frac{3}{2}}}|\Big(p_0(f,\lambda) + p_1(f,\lambda)\log(|\lambda|+1)\Big), \quad z \in \Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3}). \tag{4.34}$$

We estimate $I'_{ee}(z) = -a(z\omega)a(z\overline{\omega})V_q(z)f_e(z)$. Using (2.29), (2.30) and (2.38) we obtain

$$|I'_{ee}(z)| \leqslant C\|q\|_\infty|f_e(z)||\lambda|^{-\frac{1}{3}}\langle z\rangle^{-\frac{1}{2}}, \quad z \in \Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3}). \tag{4.35}$$

In order to estimate I_{pe} we use $P_+(s,z)$, given by (4.27). Integrating by parts we have

$$I_{pe}(z) = -P_+(z_*, z)F(z_*) + \int_{\Gamma_\lambda(z,z_*)} P_+(s,z)F'(s)\,ds, \qquad F(z) = \lambda^{-\frac{1}{6}}a(z\omega)^2 f_e(z)\rho(z).$$

Using (A.28), (2.38), (2.39), (2.29) and (2.30) we obtain

$$|I_{pe}(z)| \leqslant C\frac{\|q\|_{\mathcal{B}}}{|\lambda|^{\frac{1}{6}}}\left[|f_e(z_*)||e^{\frac{4}{3}z_*^{\frac{3}{2}}}| + \int_{\Gamma_\lambda^-}|e^{\frac{4}{3}s^{\frac{3}{2}}}|\left\{\frac{|f'_e(s)|}{\langle s\rangle^{\frac{1}{2}}} + |f_e(s)|\left(\langle s\rangle^{-\frac{3}{2}} + \frac{|\lambda|^{-\frac{2}{3}}}{\langle s\rangle^{\frac{1}{2}}}\right)\right\}|ds|\right].$$

We have $|e^{\frac{4}{3}s^{\frac{3}{2}}}| \leqslant 1$ for $s \in \Gamma_\lambda^- \subset S[-\pi, -\frac{\pi}{3})$. Using (2.46) and (4.14–4.15) we obtain

$$|I_{pe}(z)| \leqslant C\|q\|_{\mathcal{B}}|\lambda|^{-\frac{1}{6}}\Big(p_0(f,\lambda) + p_1(f,\lambda)\log(|\lambda|+1)\Big), \qquad z \in \Gamma_\lambda^-. \tag{4.36}$$

We estimate $I'_{pe}(z) = -a(z\omega)^2 e^{\frac{4}{3}z^{\frac{3}{2}}}V_q(z)f_e(z)$. Using Lemma 2.3, (2.29), (2.30) and (2.38) we get

$$\left|I'_{pe}(z)\right| \leqslant C\|q\|_\infty|e^{\frac{4}{3}z_*^{\frac{3}{2}}}f_e(z)||\lambda|^{-\frac{1}{3}}\langle z\rangle^{-\frac{1}{2}}, \qquad z \in \Gamma_\lambda^-. \tag{4.37}$$

In order to estimate I_{ep} we use $P_-(s,z)$, given by (4.27). Integrating by parts we have

$$I_{ep}(z) = -P_-(z_*, z)F(z) + \int_{\Gamma_\lambda(z,z_*)} P_-(z_*,s)F'(s)\,ds, \qquad F(z) = \lambda^{-\frac{1}{6}}a(z\overline{\omega})^2 f_p(z)\rho(z).$$

Using (A.27), (2.38), (2.39), (2.29) and (2.30) we obtain

$$|I_{ep}(z)| \leqslant C\frac{\|q\|_{\mathcal{B}}}{|\lambda|^{\frac{1}{6}}}\left[\frac{|f_p(z)e^{-\frac{4}{3}z^{\frac{3}{2}}}|}{\langle z\rangle^{\frac{1}{2}}} + \int_{\Gamma_\lambda(z,z_*)}|e^{-\frac{4}{3}s^{\frac{3}{2}}}|\left\{\frac{|f'_p(s)|}{\langle s\rangle^{\frac{1}{2}}} + |f_p(s)|\left(\langle s\rangle^{-\frac{3}{2}} + \frac{|\lambda|^{-\frac{2}{3}}}{\langle s\rangle^{\frac{1}{2}}}\right)\right\}|ds|\right].$$

By Lemma 2.3, we have $\max\limits_{\Gamma_\lambda(z,z_*)}|e^{-\frac{4}{3}s^{\frac{3}{2}}}| = |e^{-\frac{4}{3}z^{\frac{3}{2}}}|$. Using (2.46), (4.14) and (4.15) we obtain

$$|I_{ep}(z)| \leqslant C\|q\|_{\mathcal{B}}|\lambda|^{-\frac{1}{6}}|e^{-\frac{4}{3}z^{\frac{3}{2}}}|\Big(p_0(f,\lambda) + p_1(f,\lambda)\log(|\lambda|+1)\Big), \qquad z \in \Gamma_\lambda^-. \tag{4.38}$$

We estimate $I'_{ep}(z) = -a(z\overline{\omega})^2 e^{-\frac{4}{3}z^{\frac{3}{2}}} V_q(z) f_p(z)$. Using Lemma 2.3, (2.29), (2.30) and (2.38) we obtain

$$\left|I'_{ep}(z)\right| \leqslant C\|q\|_\infty |e^{-\frac{4}{3}z^{\frac{3}{2}}} f_p(z)| |\lambda|^{-\frac{1}{3}} \langle z\rangle^{-\frac{1}{2}}, \qquad z \in \Gamma_\lambda^-. \tag{4.39}$$

Due to (4.30), (4.32), (4.34), (4.36) and (4.38) we have

$$p_0(\mathbf{J}V_q f, \lambda) = |(\mathbf{j}_p V_q f)(z)| + |e^{\frac{4}{3}z^{\frac{3}{2}}} (\mathbf{j}_e V_q f)(z)| \leqslant C \frac{\|q\|_{\mathcal{B}}}{|\lambda|^{\frac{1}{6}}} \Big(p_0(f, \lambda) + p_1(f, \lambda) \log(|\lambda| + 1) \Big),$$

where $z \in \Gamma_\lambda^-$. This proves (4.28). Due to (4.30), (4.33), (4.35), (4.37) and (4.39)we have

$$p_1(\mathbf{J}V_q f, \lambda) = |\partial_z (\mathbf{j}_p V_q f)(z)| + \left| e^{\frac{4}{3}z^{\frac{3}{2}}} \partial_z (\mathbf{j}_e V_q f)(z) \right| \leqslant C \frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}} \langle z\rangle^{\frac{1}{2}}} (|f_p(z)| + |e^{\frac{4}{3}z^{\frac{3}{2}}} f_e(z)|),$$

where $z \in \Gamma_\lambda^-$. This proves (4.29). ■

Lemma 4.4. *Let* $q \in \mathcal{B}$ *and* $0 \leqslant \arg\lambda \leqslant \delta$, $|\lambda| \geqslant 1$. *Assume* $f = f^- + f^+$, *where* $f^+ = f\big|_{\Gamma_\lambda^+} \in \mathcal{F}_{\alpha,\beta}^\lambda$ *for* $\alpha > 0$, $\beta > \frac{3}{4}$ *and* $f^- = f\big|_{\Gamma_\lambda^-} \in \mathcal{F}_-^\lambda$. *Then* $g = (\mathbf{J}Vf)\big|_{\Gamma_\lambda^-} \in \mathcal{F}_-^\lambda$ *and for the decomposition* $g(z) = a(z\overline{\omega})(\mathbf{J}_p Vf)(z) + e^{\frac{4}{3}z^{\frac{3}{2}}} a(z\omega)(\mathbf{J}_e Vf)(z)$ *the following estimates are fulfilled:*

$$p_0(g, \lambda) \leqslant C \frac{\|q\|_{\mathcal{B}}}{|\lambda|^{\frac{1}{6}}} \Big(p_0(f^-, \lambda) + p_1(f^-, \lambda) \log(|\lambda| + 1) + \|f^+\|_{\alpha,\beta} \Big) + \frac{C}{|\lambda|^{\frac{2}{3}}} p_0(f^-, \lambda), \tag{4.40}$$

$$p_1(g, \lambda) \leqslant C |\lambda|^{-\frac{1}{3}} \left(\|q\|_\infty + 1\right) p_0(f^-, \lambda). \tag{4.41}$$

Proof. By (4.26), we have $g = g_0 + g_- + g_+$, where

$$g_+(z) = a(z\overline{\omega}) h_p(V_q f^+) + e^{\frac{4}{3}z^{\frac{3}{2}}} a(z\omega) h_e(V_q f^+),$$

$$g_-(z) = a(z\overline{\omega}) (\mathbf{j}_p V_q f^-)(z) + e^{\frac{4}{3}z^{\frac{3}{2}}} a(z\omega) (\mathbf{j}_e V_q f^-)(z),$$

$$g_0(z) = a(z\overline{\omega}) (\mathbf{J}_p V_0 f)(z) + e^{\frac{4}{3}z^{\frac{3}{2}}} a(z\omega) (\mathbf{J}_e V_0 f)(z).$$

Firstly, we estimate g_+. From (3.15) and (3.25) we have

$$|h_p(V_q f)| \leqslant C \|q_1\|_\infty |\lambda|^{-\frac{1}{6}} \|f^+\|_{\alpha,\beta}, \quad |h_e(V_q f)| \leqslant C(\|q\|_\infty |e^{-\frac{4}{3}z_*^{\frac{3}{2}}}| + \|q_1\|_\infty) |\lambda|^{-\frac{1}{6}} \|f^+\|_{\alpha,\beta}. \tag{4.42}$$

By Lemma 2.3, we have $|e^{\frac{4}{3}(z^{\frac{3}{2}} - z_*^{\frac{3}{2}})}| \leqslant 1$ for $z \in \Gamma_\lambda^-$. Thus

$$p_0(g_+, \lambda) \leqslant C(\|q\|_\infty + \|q_1\|_\infty) |\lambda|^{-\frac{1}{6}} \|f^+\|_{\alpha,\beta}, \qquad p_1(g_+, \lambda) = 0. \tag{4.43}$$

Secondly, we estimate g_0. Using (4.21), (4.24), (2.29), (2.30), (2.38), (2.47) and Lemma 2.3 gives

$$|(\mathbf{j}_p V_0 f)(z)| \leqslant C \int_{\Gamma_\lambda^-} \frac{p_0(f^-, \lambda)}{|\lambda|^{\frac{4}{3}} + |s|^2} \frac{|ds|}{\langle s\rangle^{\frac{1}{2}}} \leqslant C \frac{p_0(f^-, \lambda)}{|\lambda|^{\frac{2}{3}}}, \qquad z \in \Gamma_\lambda^-, \tag{4.44}$$

$$|(\mathbf{j}_e V_0 f)(z)| \leqslant C|e^{-\frac{4}{3}z^{\frac{3}{2}}}|\int_{\Gamma_\lambda^-} \frac{p_0(f^-,\lambda)}{|\lambda|^{\frac{4}{3}}+|s|^2}\frac{|ds|}{\langle s\rangle^{\frac{1}{2}}} \leqslant C|e^{-\frac{4}{3}z^{\frac{3}{2}}}|\frac{p_0(f^-,\lambda)}{|\lambda|^{\frac{2}{3}}} \qquad z\in\Gamma_\lambda^-. \tag{4.45}$$

Using Lemma 2.3, (3.29) and (3.30) we have

$$|h_p(V_0 f)| \leqslant C|\lambda|^{-\frac{2}{3}}\|f^+\|_\alpha, \qquad |h_e(V_0 f)| \leqslant C|\lambda|^{-\frac{2}{3}}|e^{-\frac{4}{3}z_*^{\frac{3}{2}}}|\|f^+\|_\alpha. \tag{4.46}$$

By Lemma 2.3, we have $|e^{\frac{4}{3}(z^{\frac{3}{2}}-z_*^{\frac{3}{2}})}| \leqslant 1$ for $z\in\Gamma_\lambda^-$. Thus substituting (4.44), (4.45) and (4.46) in (4.26) we obtain

$$p_0(g_0,\lambda) \leqslant C(p_0(f^-,\lambda)+\|f^+\|_\alpha)|\lambda|^{-\frac{2}{3}}. \tag{4.47}$$

Using (4.21), (4.24), (2.38), (2.29), (2.30) and (4.13) we obtain

$$|\partial_z(\mathbf{j}_p V_0 f)(z)| \leqslant \frac{C}{|\lambda|^{\frac{4}{3}}}\frac{p_0(f^-,\lambda)}{\langle z\rangle^{\frac{1}{2}}}, \qquad |\partial_z(\mathbf{j}_e V_0 f)(z)| \leqslant \frac{C}{|\lambda|^{\frac{4}{3}}}|e^{-\frac{4}{3}z^{\frac{3}{2}}}|\frac{p_0(f^-,\lambda)}{\langle z\rangle^{\frac{1}{2}}},$$

which proves

$$p_1(g_0,\lambda) \leqslant Cp_0(f^-,\lambda)|\lambda|^{-\frac{4}{3}}. \tag{4.48}$$

Finally, apply (4.28) to g_-; together with (4.43) and (4.47) this gives (4.40). Applying (4.29) to g_- together with (4.43) and (4.48) gives (4.41). ■

Proof of Theorem 4.1. We consider the case $0\leqslant \arg\lambda\leqslant\delta$, the proof for $-\delta\leqslant\arg\lambda\leqslant 0$ is similar. Let $v_{n+1}=\mathbf{J}Vv_n, n\geqslant 0$, where $v_0\equiv a$. Introduce $v_n^\pm = v_n\big|_{\Gamma_\lambda^\pm}$. By (3.31–3.32), for some absolute constant $c_0>0$ we have

$$\|v_{n+1}^+\|_{\alpha_{n+1},\beta_{n+1}} \leqslant \varepsilon\|v_n^+\|_{\alpha_n,\beta_n},$$

where α_n and β_n are given by (3.33). We estimate v_n^- in terms of p_0 and p_1, using the decomposition (4.16) for v_0^- and (4.18), (4.19) for v_n^-, $n\geqslant 1$. Substituting $f=v_n$ and $g=v_{n+1}$ in (4.40), (4.41) and choosing c_0 sufficiently large, we obtain

$$p_0(v_{n+1}^-,\lambda) \leqslant \varepsilon\Big(p_0(v_n^-,\lambda)+p_1(v_n^-,\lambda)\log(|\lambda|+1)+\|v_n^+\|_{\alpha_n,\beta_n}\Big), \quad p_1(v_{n+1}^-) \leqslant \varepsilon\frac{p_0(v_n^-,\lambda)}{|\lambda|^{\frac{1}{6}}}.$$

Using (3.8) and (4.17) (in particular, $p_1(v_0,\lambda)=0$) we obtain for v_1 and v_2

$$p_0(v_1^-,\lambda)\leqslant\varepsilon L, \qquad p_1(v_1^-,\lambda)\leqslant\varepsilon|\lambda|^{-\frac{1}{6}}L, \qquad L=2+\|v_0\|_{\frac{1}{4},\frac{5}{4}},$$

$$p_0(v_2^-,\lambda)\leqslant\varepsilon\Big(\varepsilon L+\varepsilon L|\lambda|^{-\frac{1}{6}}\log(|\lambda|+1)+\|v_1^+\|_{\alpha_1,\beta_1}\Big)\leqslant\varepsilon^2(2+|\lambda|^{-\frac{1}{6}}\log(|\lambda|+1))L,$$

$$p_1(v_2^-,\lambda)\leqslant\varepsilon^2|\lambda|^{-\frac{1}{6}}L.$$

Increasing the constant c_0 and using the induction principle we obtain for each integer $n \geqslant 0$

$$p_0(v_n^-, \lambda) \leqslant \varepsilon^n L, \quad p_1(v_n^-, \lambda) \leqslant \frac{1}{|\lambda|^{\frac{1}{6}}} \varepsilon^n L, \quad \|v_n^+\|_{\alpha_n, \beta_n} \leqslant \varepsilon^n \|v_0\|_{\frac{1}{4}, \frac{5}{4}}. \tag{4.49}$$

By Theorem 3.1, for $\varepsilon < 1$ the series $v_+\Big|_{\Gamma_\lambda^+} = \sum_{n=0}^\infty v_n^+$ converges in $\mathcal{F}_{\frac{1}{4},1}$-norm and gives a solution of $v_+ = v_0 + \mathbf{J}Vv_+$ on Γ_λ^+. By (4.49), for $\varepsilon < 1$ the series $\sum_{n=0}^\infty p_0(v_n^-, \lambda)$ and $\sum_{n=0}^\infty p_1(v_n^-, \lambda)$ converge; for $\varepsilon \leqslant \frac{1}{2}$ the following estimates are fulfilled:

$$p_0(v_+|_{\Gamma_\lambda^-}, \lambda) \leqslant 2L, \quad p_1(v_+|_{\Gamma_\lambda^-}, \lambda) \leqslant \frac{2L}{|\lambda|^{\frac{1}{6}}}, \quad p_0(v_1^-, \lambda) \leqslant 2\varepsilon L, \quad p_1(v_1^-, \lambda) \leqslant \frac{2\varepsilon L}{|\lambda|^{\frac{1}{6}}}, \tag{4.50}$$

$$p_0\left(v_+|_{\Gamma_\lambda^-} - v_0^- - v_1^-, \lambda\right) \leqslant 2\varepsilon^2 L, \qquad p_1\left(v_+|_{\Gamma_\lambda^-} - v_0^- - v_1^-, \lambda\right) \leqslant 2|\lambda|^{-\frac{1}{6}} \varepsilon^2 L. \tag{4.51}$$

By (4.13), convergence of $\sum_{n=0}^\infty p_0(v_n^-, \lambda)$ and $\sum_{n=0}^\infty p_1(v_n^-, \lambda)$ implies convergence of $\sum_{n=0}^\infty v_n^-$ in C^1-norm on Γ_λ^-. Thus for $\varepsilon < 1$ the series $v_+ = \sum_{n=0}^\infty v_n$ converges and solves the equation $v_+ = v_0 + \mathbf{J}Vv_+$ on Γ_λ. Using (4.13) and (4.49) we obtain (4.1) and (4.2) from (4.50); similarly we obtain (4.3) from (4.51).

We prove uniqueness. Suppose that there exists another solution $v_+^{(1)} \in \mathcal{F}^\lambda = \mathcal{F}_-^\lambda \oplus \mathcal{F}_{\frac{1}{4},1}^\lambda$. By Theorem 3.1, we have $v_+^{(1)}|_{\Gamma_\lambda^+} = v_+|_{\Gamma_\lambda^+}$. Consider the difference $y = (v_+ - v_+^{(1)})|_{\Gamma_\lambda^-} \in \mathcal{F}_-^\lambda$. We have $y = (\mathbf{J}V)^n y$ for any $n > 1$; applying Lemma 4.4 we obtain $p_0(y, \lambda) \leqslant C\varepsilon^n(p_0(y, \lambda) + p_1(y, \lambda))$. Taking the limit $n \to \infty$ for $\varepsilon < 1$ gives $y = 0$. ■

5 Asymptotics of the Wronskian

In this section we shall determine the asymptotics of $w(\lambda) = \{\psi_-, \psi_+\}$ as $\lambda \to \infty$. To this end we find the asymptotics of $u_1(z, \lambda) = e^{-\frac{2}{3}z^{\frac{3}{2}}}(\mathbf{J}Vv_0)(z)$ (given by Corollary 4.2) as $\lambda \to \infty$ in the sector $|\arg \lambda| \leqslant \delta$. Introduce an auxiliary function

$$E_+(\lambda) = 2\int_{\Gamma_\lambda^-} \mathrm{Ai}(s\omega)\mathrm{Ai}(s\overline{\omega})V_q(s)\, ds, \qquad \omega = e^{\frac{2\pi i}{3}}. \tag{5.1}$$

By (2.27), we have

$$\frac{2}{3} z_0^{\frac{3}{2}} = \pm i \frac{\pi}{4}\lambda, \qquad 0 \leqslant \pm \arg \lambda \leqslant \pi, \qquad z_0 = z(0, \lambda) = -\lambda^{\frac{2}{3}} \left(\frac{3\pi}{8}\right)^{\frac{2}{3}}. \tag{5.2}$$

Lemma 5.1. *1. Let $q \in \mathcal{B}$ and $|\arg \lambda| \leqslant \delta$, $|\lambda| \to \infty$. Then*

$$u_+(z_0, \lambda) = z_0^{-\frac{1}{4}} \left[\sin\frac{\pi}{4}(\lambda + 1)\left(1 + O(\lambda^{-\frac{1}{6}})\right) + E_+(\lambda)\cos\frac{\pi}{4}(\lambda + 1)\right] + O\left(\frac{e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}}{\lambda^{\frac{1}{2}}}\right), \tag{5.3}$$

$$\partial_z u_+(z_0,\lambda) = z_0^{\frac{1}{4}}\left[-\cos\frac{\pi}{4}(\lambda+1)\left(1+O(\lambda^{-\frac{1}{6}})\right) + E_+(\lambda)\sin\frac{\pi}{4}(\lambda+1)\right] + O\left(\frac{e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}}{\lambda^{\frac{1}{6}}}\right), \tag{5.4}$$

2. Let $q \in \mathcal{B}$ *and* $\delta \leqslant \pm\arg\lambda \leqslant \pi$, $|\lambda| \to \infty$. *Then*

$$u_+(z_0,\lambda) = \frac{e^{\mp i\frac{\pi}{4}(\lambda+1)}}{2z_0^{\frac{1}{4}}} + O\left(\frac{e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}}{\lambda^{\frac{2}{3}}}\right), \quad \partial_z u_+(z_0,\lambda) = -\frac{z_0^{\frac{1}{4}}}{2}e^{\mp i\frac{\pi}{4}(\lambda+1)} + O\left(\frac{e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}}{\lambda^{\frac{1}{3}}}\right). \tag{5.5}$$

Proof. We present the proof only for $\operatorname{Im}\lambda \geqslant 0$, for $\operatorname{Im}\lambda \leqslant 0$ it is analogous.

1. Let $0 \leqslant \arg\lambda \leqslant \delta$. We have

$$u_1(z,\lambda) = e^{-\frac{2}{3}z^{\frac{3}{2}}}v_1(z) = e^{-\frac{2}{3}z^{\frac{3}{2}}}(\mathbf{J}Vv_0)(z) = \operatorname{Ai}(z\overline{\omega})(\mathbf{J}_pVv_0)(z) + \operatorname{Ai}(z\omega)(\mathbf{J}_eVv_0)(z).$$

By (4.47), $(\mathbf{J}V_0v_0)(z) = O(\lambda^{-\frac{2}{3}})$. Since $V = V_q + V_0$, we have as $\lambda \to \infty$

$$u_1(z,\lambda) = e^{-\frac{2}{3}z^{\frac{3}{2}}}v_1(z) = e^{-\frac{2}{3}z^{\frac{3}{2}}}(\mathbf{J}V_qv_0)(z) + O\left(|\lambda|^{-\frac{2}{3}}\langle z\rangle^{-\frac{1}{4}}e^{\frac{2}{3}|\operatorname{Re} z^{\frac{3}{2}}|}\right). \tag{5.6}$$

Using (4.16), (2.32), (2.33), (3.6), (4.9) and (4.20–4.25) we obtain

$$\begin{gathered}
e^{-\frac{2}{3}z^{\frac{3}{2}}}(\mathbf{J}V_qv_0)(z) = \operatorname{Ai}(z\overline{\omega})(\mathbf{J}_pV_qv_0)(z) + \operatorname{Ai}(z\omega)(\mathbf{J}_eV_qv_0)(z) \\
= 2i\omega\operatorname{Ai}(z)\int_{\Gamma_\lambda^+} a(s\omega)a(s)V_q(s)\,ds - 2i\operatorname{Ai}(z\omega)\int_{\Gamma_\lambda^+} e^{-\frac{4}{3}s^{\frac{3}{2}}}a^2(s)V_q(s)\,ds \\
+2\operatorname{Bi}(z)\int_{\Gamma_\lambda(z,z_*)} a(s\omega)a(s\overline{\omega})V_q(s)\,ds - 2i\overline{\omega}\operatorname{Ai}(z\overline{\omega})\int_{\Gamma_\lambda(z,z_*)} e^{\frac{4}{3}s^{\frac{3}{2}}}a^2(s\omega)V_q(s)\,ds \\
+2ie^{i\frac{\pi}{3}}\operatorname{Ai}(z\omega)\int_{\Gamma_\lambda(z,z_*)} e^{-\frac{4}{3}s^{\frac{3}{2}}}a^2(s\overline{\omega})V_q(s)\,ds,
\end{gathered}$$

where $\omega = e^{\frac{2\pi i}{3}}$. Now we set $z = z_0$ and write the asymptotics of u_1 in terms of E_+. By Lemma 7.7, (3.6), (4.9) and (2.38), we have

$$u_1(z_0,\lambda) = 2i\omega\operatorname{Ai}(z_0)\int_{\Gamma_\lambda^+} a(s\omega)a(s)V_q(s)\,ds + \operatorname{Bi}(z_0)E_+(\lambda) + O\Big(\lambda^{-\frac{1}{3}}z_0^{-\frac{1}{4}}|e^{-\frac{2}{3}z_0^{\frac{3}{2}}}|\Big). \tag{5.7}$$

Using (2.39) we obtain for the derivative

$$\partial_z u_1(z,\lambda) = 2i\omega\operatorname{Ai}'(z_0)\int_{\Gamma_\lambda^+} a(s\omega)a(s)V_q(s)\,ds + \operatorname{Bi}'(z_0)E_+(\lambda) + O\Big(\lambda^{-\frac{1}{3}}z^{\frac{1}{4}}|e^{-\frac{2}{3}z^{\frac{3}{2}}}|\Big). \tag{5.8}$$

From Lemma 3.2, (3.6), (2.38) and (2.39) we deduce that $\int_{\Gamma_\lambda^+} a(s\omega)a(s)V_q(s)\,ds = O(\lambda^{-\frac{1}{6}})$. Therefore using (4.6), (4.7), (5.2), (5.6), (5.7) and (5.8) we obtain

$$u_+(z_0,\lambda) = \operatorname{Ai}(z_0)(1+O(\lambda^{-\frac{1}{6}})) + \operatorname{Bi}(z_0)E_+(\lambda) + O\left(\lambda^{-\frac{1}{2}}e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}\right), \tag{5.9}$$

$$\partial_z u_+(z_0,\lambda) = \mathrm{Ai}'(z_0)(1+O(\lambda^{-\frac16})) + \mathrm{Bi}'(z_0)E_+(\lambda) + O\left(\lambda^{-\frac16} e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}\right), \qquad (5.10)$$

where z_0 is given by (2.27). Recall the standard uniform asymptotics of Airy functions [1] in the sector $|\arg z| < \frac{\pi}{3} - \varepsilon$, $\varepsilon > 0$, as $z \to \infty$:

$$\mathrm{Ai}(-z) = z^{-\frac14}(\sin\eta + O(F(z)), \quad \mathrm{Ai}'(-z) = -z^{\frac14}(\cos\eta + O(F(z))), \qquad (5.11)$$

$$\mathrm{Bi}(-z) = z^{-\frac14}(\cos\eta + O(F(z)), \quad \mathrm{Bi}'(-z) = z^{\frac14}(\sin\eta + O(F(z))), \qquad (5.12)$$

where $\eta = \frac23 z^{\frac32} + \frac{\pi}{4}$ and $F(z) = z^{-\frac32} e^{\frac23 |\operatorname{Im} z^{\frac32}|}$. Note that for $0 \leqslant \arg\lambda \leqslant \delta$ we have $-\pi \leqslant \arg z_0 \leqslant -\pi + \frac23\delta$. Thus we substitute (5.11) and (5.12) into (5.9–5.10). This gives (5.3),(5.4).

2. Let $\delta \leqslant \arg\lambda \leqslant \pi$. By (4.8), we have

$$u_+(z_0,\lambda) = \mathrm{Ai}(z_0) + O\left(\lambda^{-\frac23} e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}\right), \quad u_+'(z_0,\lambda) = \mathrm{Ai}'(z_0) + O\left(\lambda^{-\frac13} e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}\right). \qquad (5.13)$$

Note that for $\delta \leqslant \arg\lambda \leqslant \pi$ we have $-\pi + \frac23\delta \leqslant \arg z_0 \leqslant 0$. Thus we apply (2.31) and (5.2) to (5.13), which gives (5.5).■

Introduce the function

$$\phi(\lambda) = 2^{\frac34}\left(\frac{\lambda}{2e}\right)^{\lambda/4}, \quad \lambda \in \mathbb{C}\setminus\mathbb{R}_-, \qquad \phi(\lambda) > 0 \quad \text{for} \quad \lambda > 0. \qquad (5.14)$$

In the next Lemma we write ψ_+, defined by (2.2), in terms of u_+.

Lemma 5.2. *Let $q \in \mathcal{B}$. Let ψ_+ be the solution of Eq.(2.1), satisfying (2.2); let u_+ be the solution of Eq.(2.34), given by Corollary 4.2. Then for $|\lambda|^{\frac16} \geqslant c_0\left(\|q\|_{\mathcal{B}} + 1\right)$, where $c_0 \geqslant 1$ is some absolute constant, the following identity holds:*

$$\psi_+(x,\lambda) = \phi_+(x,\lambda), \quad \textit{where} \quad \phi_+(x,\lambda) = \phi(\lambda)\frac{u_+(z(x,\lambda)),\lambda)}{\sqrt{z'(x,\lambda)}}, \qquad x \geqslant 0. \qquad (5.15)$$

Proof. The function ϕ_+, given by (5.15), solves Eq.(2.1) by changing variables according to (2.26). In order to prove (5.15) it is sufficient to demonstrate that ϕ_+ has the asymptotics (2.2). Using (2.21), (2.26), (2.31), Corollary 4.2 and (5.2), we have for $x \to \infty$

$$\phi_+(x,\lambda) = e^{-\frac{x^2}{2}}(\sqrt{2}x)^{\frac{\lambda-1}{2}}\Big(1 + O\Big(\frac{1}{x^{\frac13}}\Big)\Big), \quad \frac{\partial\phi_+(x,\lambda)}{\partial x} = -\frac{e^{-\frac{x^2}{2}}}{\sqrt{2}}(\sqrt{2}x)^{\frac{\lambda+1}{2}}\Big(1 + O\Big(\frac{1}{x^{\frac13}}\Big)\Big),$$

which proves (5.15). ■

In order to obtain the asymptotics of the Wronskian $w(\lambda) = \{\psi_-, \psi_+\}$ we need also the asymptotics of the fundamental solution ψ_- (see Theorem 2.1). We use our results for $x > 0$ and consider the reflected potential $q_-(x) = q(-x)$ for $x \in \mathbb{R}_+$. We define

$$V_q^-(z) = \lambda^{-\frac13}\rho^2(z,\lambda)\hat{q}_-(z), \qquad \hat{q}_-(z,\lambda) \equiv q_-(\sqrt{\lambda}t(z\lambda^{-\frac23})),$$

where ρ is given by (2.29). Let $u_-(z,\lambda) = u_+(z,\lambda;V_q^-)$, where u_+ is given by Corollary 4.2 (with V_q replaced by V_q^-). By Lemma 5.2, the fundamental solution ψ_- is related to u_- by

$$\psi_-(-x,\lambda) = \phi(\lambda)\frac{u_-(z(x,\lambda),\lambda)}{\sqrt{z'(x,\lambda)}}, \qquad x \geqslant 0. \tag{5.16}$$

Introduce $E_-(\lambda)$ by

$$E_-(\lambda) = 2\int_{\Gamma_\lambda^-} \mathrm{Ai}(s\omega)\mathrm{Ai}(s\overline{\omega})V_q^-(s)\,ds, \qquad \omega = e^{\frac{2\pi i}{3}}. \tag{5.17}$$

By Lemma 5.1, for $|\arg\lambda| \geqslant \delta$ the solution $u_-(z_0,\lambda)$ has the asymptotics (5.5); for $|\arg\lambda| \leqslant \delta$ it has the asymptotics (5.3), (5.4) with E_+ replaced by E_-:

$$u_-(z_0,\lambda) = z_0^{-\frac{1}{4}}\left[\sin\frac{\pi}{4}(\lambda+1)\left(1+O(\lambda^{-\frac{1}{6}})\right) + E_-(\lambda)\cos\frac{\pi}{4}(\lambda+1)\right] + O\left(\lambda^{-\frac{1}{2}}e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}\right), \tag{5.18}$$

$$u_-'(z_0,\lambda) = z_0^{\frac{1}{4}}\left[-\cos\frac{\pi}{4}(\lambda+1)\left(1+O(\lambda^{-\frac{1}{6}})\right) + E_-(\lambda)\sin\frac{\pi}{4}(\lambda+1)\right] + O\left(\lambda^{-\frac{1}{6}}e^{\frac{\pi}{4}|\operatorname{Im}\lambda|}\right). \tag{5.19}$$

Below we use the function $E = E_- + E_+$.

Lemma 5.3. *Let $q \in \mathcal{B}$. Then for $|\lambda| \to \infty$ the following uniform asymptotics hold:*

$$w(\lambda) = \phi^2(\lambda)\left[\cos\frac{\pi}{2}\lambda - E(\lambda)\sin\frac{\pi}{2}\lambda + O\left(\lambda^{-\frac{1}{3}}e^{\frac{\pi}{2}|\operatorname{Im}\lambda|}\right)\right], \qquad |\arg\lambda| \leqslant \delta \tag{5.20}$$

$$w(\lambda) = \frac{\phi^2(\lambda)}{2}\left[e^{\mp i\frac{\pi}{2}\lambda} + O\left(\frac{e^{\frac{\pi}{2}|\operatorname{Im}\lambda|}}{\lambda^{\frac{1}{6}}}\right)\right] = \phi^{-2}(-\lambda)\left(4+O(\lambda^{-\frac{1}{6}})\right), \qquad \delta \leqslant \pm\arg\lambda \leqslant \pi. \tag{5.21}$$

Proof. Using (2.26), (5.15) and (5.16) we obtain

$$w(\lambda) = -\phi^2(\lambda)\Big(u_+(z_0,\lambda)u_-'(z_0,\lambda) + u_+'(z_0,\lambda)u_-(z_0,\lambda)\Big) - \phi^2(\lambda)\frac{u_+(z_0,\lambda)u_-(z_0,\lambda)}{16\lambda k^2(0)}.$$

Substituting (5.3–5.4) and (5.18–5.19) in the last identity, we obtain (5.20); substituting (5.5) and the same formulae for u_-, we obtain (5.21). ■

6 The proof of Theorem 1.1

Using (5.14) we write the following uniform asymptotics of the unperturbed Wronskian $w_0(\lambda) = \{\psi_-^0, \psi_+^0\} = -\frac{2\sqrt{\pi}}{\Gamma(\frac{1-\lambda}{2})}$ (see, for example, [1]):

$$w_0(\lambda) = \phi^2(\lambda)\cos\frac{\pi\lambda}{2}\cdot\left(1+O\left(\lambda^{-1}\right)\right), \qquad |\arg\lambda| \leqslant \delta \tag{6.1}$$

$$w_0(\lambda) = \phi^{-2}(-\lambda)\left(4+O\left(\lambda^{-1}\right)\right), \qquad |\arg\lambda| \geqslant \delta. \tag{6.2}$$

Lemma 6.1. *Let $q \in \mathcal{B}$. Then there is $N_0 \in \mathbb{Z}$ such that for each integer $N > N_0$ the operator T has exactly N simple eigenvalues in the disc $\{z : |z| < 2N\}$ and for each $n > N$, exactly one simple eigenvalue in the disc $\{z : |z - \mu_n^0| < n^{-\frac{1}{6}}\}$. There are no other eigenvalues.*

Proof. Consider the contours $|\lambda| = 2n$, $|\lambda - \mu_n^0| = \delta_n$, $n > N$, $n \in \mathbb{N}$, where $\delta_n = n^{-\frac{1}{6}} < \frac{\log 2}{\pi}$. Then $|\cos\frac{\pi\lambda}{2}| \geqslant 4e^{\frac{\pi}{2}|\mathrm{Im}\,\lambda|}$ for $|\lambda| = 2n$ and $\left|\frac{1}{\pi}\cos\frac{\pi\lambda}{2}\right| \geqslant \frac{\delta}{4}e^{\frac{\pi}{2}|\mathrm{Im}\,\lambda|}$ for $|\lambda - \mu_n^0| = \delta$, $\delta < \frac{\log 2}{\pi}$. By the asymptotics (5.20), (5.21), (6.2) and (6.1), there exist integer $N_0 > \left(\frac{\pi}{\log 2}\right)^6$ such that for any integer $N > N_0$ on these contours $|w(\lambda) - w_0(\lambda)| \leqslant \frac{1}{2}|w_0(\lambda)|$. It follows that $w(\lambda)$ does not vanish on these contours. Hence, by Rouche's theorem, $w(\lambda)$ has as many roots, counted with multiplicities, as $w_0(\lambda)$ in each of the bounded regions and in the remaining unbounded region. Since $w_0(\lambda)$ has only the simple roots $\{\mu_n^0\}_{n=0}^{\infty}$, the Lemma is proved. ■

Lemma 6.2. *Let $q \in \mathcal{B}$. Then for $|\lambda| \to \infty$ the following asymptotics are fulfilled:*

$$E(\lambda + \varepsilon) - E(\lambda) = O\left(\lambda^{-\frac{1}{3}}\right), \qquad \varepsilon = O\left(\lambda^{-\frac{1}{6}}\right), \qquad -\frac{\delta}{2} \leqslant \arg\lambda \leqslant \frac{\delta}{2}, \tag{6.3}$$

$$E(\lambda) = -\frac{1}{2}\int_{-1}^{1}\frac{q(t\sqrt{\lambda})dt}{\sqrt{1-t^2}} + O\left(\|q\|_{\mathcal{B}}\lambda^{-\frac{1}{3}}\right), \qquad E(\lambda) = O\left(\|q\|_{\mathcal{B}}\lambda^{-\frac{1}{4}}\right), \qquad \lambda > 0. \tag{6.4}$$

Proof. Consider the case $0 \leqslant \arg\lambda \leqslant \delta$; for $-\delta \leqslant \arg\lambda \leqslant 0$ the proof is analogous. Consider E_+, given by (5.1). Recall that $t_* = \frac{x_*}{\sqrt{\lambda}}$, where x_* is defined by $z_* = z(x_*, \lambda)$ (see Lemma 2.2 and below). We have $\lambda = |\lambda|e^{2i\vartheta}$. For some $c \in [0, 1]$ we set $t_1 = t_* - \frac{c}{|\lambda|^{\frac{2}{3}}}e^{-i\vartheta}$, $x_1 = \frac{t_1}{\sqrt{\lambda}}$ and $z_1 = z(x_1, \lambda)$. The length of $\Gamma_\lambda(z_1, z_*)$ is $|\Gamma_\lambda(z_1, z_*)| = |\lambda|^{\frac{2}{3}}\int_{[t_1, t_*]}|k'(t)|\,|dt|$. By Lemma 7.3.4 , $|t_*|$ is bounded uniformly in λ. Thus using (2.21) we conclude that $|\Gamma_\lambda(z_1, z_*)| \leqslant C$. Therefore using (2.29), (2.30), (2.38), (3.6) and (4.9) gives

$$\left|\int_{\Gamma_\lambda(z_1, z_*)}\mathrm{Ai}(s\overline{\omega})\mathrm{Ai}(s\omega)V_q(s)\,ds\right| \leqslant C \cdot \frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}}. \tag{6.5}$$

Next, using the asymptotics (2.31) and Lemma 7.5.1 , we have uniformly in $|\arg\lambda| \leqslant \delta$

$$\left|\mathrm{Ai}(z\omega)\mathrm{Ai}(z\overline{\omega}) + \frac{i}{4\pi z^{\frac{1}{2}}}\right| \leqslant \frac{C}{|z|^{\frac{1}{2}}(1 + |z|)^{\frac{3}{2}}}, \qquad z \in \Gamma_\lambda^- \subset S[\pi, -\frac{\pi}{3}]. \tag{6.6}$$

Substituting (6.5) and (6.6) in (5.1), we have

$$E_+(\lambda) = 2\pi\int_{\Gamma_\lambda(z_0, z_1)}\mathrm{Ai}(s\omega)\mathrm{Ai}(s\overline{\omega})V_q(s)\,ds + O\left(\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}}\right) = -\frac{i}{2}\int_{\Gamma_\lambda(z_0, z_1)}V_q(s)s^{-\frac{1}{2}}\,ds + O\left(\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}}\right). \tag{6.7}$$

Making the substitution $s = k(t) \cdot \lambda^{\frac{2}{3}}$, $ds = \lambda^{\frac{2}{3}}k'(t)\,dt$ and using (2.29) we obtain

$$E_+(\lambda) = -\frac{i}{2}\int_{e^{-i\vartheta}[0, t_1]}\frac{q(\sqrt{\lambda}t)}{\sqrt{k(t)}k'(t)}\,dt + O\left(\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}}\right). \tag{6.8}$$

Using (2.17) and (2.19) we have $k'(t)\sqrt{k(t)} = \sqrt{t^2-1} = i\sqrt{1-t^2}$, where the last root is positive for $-1 < t^2 < 1$. Hence

$$E_+(\lambda) = -\frac{1}{2}\int_0^d \frac{q(t|\lambda|^{\frac{1}{2}})}{\sqrt{e^{2i\vartheta}-t^2}}\,dt + O\left(\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}}\right), \qquad d = 1 - \frac{c}{|\lambda|^{\frac{2}{3}}}.$$

Using similar arguments for E_-, we obtain

$$E(\lambda) = E_c(\lambda) + O\left(\frac{\|q\|_{\mathcal{B}}}{\lambda^{\frac{1}{3}}}\right), \qquad E_c(\lambda) = -\frac{1}{2}\int_{-d}^d \frac{q(t\sqrt{|\lambda|})dt}{\sqrt{e^{2i\vartheta}-t^2}}. \tag{6.9}$$

In order to prove (6.3) we set $c > 0$. We estimate the partial derivatives of $E_c(\lambda)$ with respect to real and complex parts of $\lambda = \mu + i\nu = |\lambda|e^{2i\vartheta}$. This gives $|\frac{\partial}{\partial\mu}E_c(\lambda)| \leqslant C\frac{\|q\|_{\mathcal{B}}}{|\lambda|^{\frac{2}{3}}}$, $|\frac{\partial}{\partial\nu}E_c(\lambda)| \leqslant C\frac{\|q\|_{\mathcal{B}}}{|\lambda|^{\frac{2}{3}}}$. Therefore as $\lambda \to \infty$

$$|E_c(\lambda+\varepsilon) - E_c(\lambda)| \leqslant |\varepsilon| \cdot \sup_{|\lambda-\lambda'|\leqslant\varepsilon} |\nabla E_c(\lambda')| \leqslant C\|q\|_{\mathcal{B}}|\lambda|^{-\frac{5}{6}}, \qquad \varepsilon = O\left(|\lambda|^{-\frac{1}{6}}\right),$$

which together with (6.9) proves (6.3).

Setting $c = 0$ and $\lambda > 0$ in (6.9) proves the first asymptotics in (6.4). In order to prove the second one we use the decomposition $E_0(\lambda) = E_1(\lambda) + I_2(\lambda)$, $\lambda > 0$, where

$$I_2(\lambda) = -\frac{1}{2}\int_{1-\lambda^{-\frac{1}{2}}}^1 \frac{q(t\sqrt{\lambda}) + q(-t\sqrt{\lambda})}{\sqrt{1-t^2}}dt.$$

Using $\int_{\lambda^{\frac{1}{2}}t}^{\lambda^{\frac{1}{2}}-1} q(x)\,dx = q_1(\lambda^{\frac{1}{2}}-1) - q_1(\lambda^{\frac{1}{2}}t)$ we integrate the expression for E_1 by parts. This gives $|E_1(\lambda)| \leqslant C\frac{\|q_1\|_\infty}{\lambda^{\frac{1}{4}}}$. Direct estimate gives $|I_2(\lambda)| \leqslant \frac{\|q\|_\infty}{2\lambda^{\frac{1}{4}}}$. Combining these estimates proves the second relation in (6.4). ■

Proof of Theorem 1.1 By Lemma 6.1, in each disc $D_n = \{\lambda : |\lambda - \mu_n^0| \leqslant \frac{1}{n^{\frac{1}{6}}}\}$ there exists exactly one simple eigenvalue μ_n for n sufficiently large; now we improve this estimate. Using the asymptotics (5.20) and $|\sin(\frac{\pi}{2}\lambda)| \geqslant \frac{1}{2}$ for $\lambda \in D_n$, we obtain

$$\cot(\frac{\pi}{2}\mu_n) + E(\mu_n) + O\left(n^{-\frac{1}{3}}\right) = 0, \qquad n \to \infty. \tag{6.10}$$

Using (6.4), we write

$$\mu_n - \mu_n^0 = -\frac{2}{\pi}E(\mu_n^0) + \varepsilon_n, \qquad \varepsilon_n = O(n^{-\frac{1}{6}}), \qquad n \to \infty. \tag{6.11}$$

Substituting (6.11) into (6.10) we have

$$\frac{\pi}{2}\varepsilon_n + E(\mu_n) - E(\mu_n^0) = O\left(n^{-\frac{1}{3}}\right). \tag{6.12}$$

By (6.3), $E(\mu_n) - E(\mu_n^0) = O(n^{-\frac{1}{3}})$. Thus in (6.11) the error term ε_n is $O(n^{-\frac{1}{3}})$, which yields $\mu_n = \mu_n^0 - \frac{2}{\pi}E(\mu_n^0) + O(n^{-\frac{1}{3}})$. The change of variable $t = \sin\theta$ gives $\mu_n^1 = -\frac{2}{\pi}E(\mu_n^0) = (2\pi)^{-1}\int_{-\pi}^{\pi} q(\sqrt{\mu_n^0}\sin\theta)\,d\theta$. This proves (1.2). ■

Proof of Proposition 1.2. Substituting $q(x) = \int_{\mathbb{R}} e^{ixt}\,d\nu(t)$ into (6.9) and using the identity for Bessel function $J_0(z) = \frac{1}{\pi}\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{iz\sin\phi}\,d\phi$ (see [1]) we have

$$\frac{1}{2\pi}\int_{-\pi}^{\pi} q(\sqrt{\lambda}\sin\vartheta)\,d\vartheta = \int_{\mathbb{R}} J_0(t\sqrt{\lambda})\,d\nu(t) = I_1 + I_2, \qquad \lambda > 0, \tag{6.13}$$

where

$$I_1 = \int_{|t|<\varepsilon} J_0(t\sqrt{\lambda})\,d\nu(t), \qquad I_2 = \int_{|t|>\varepsilon} J_0(t\sqrt{\lambda})\,d\nu(t), \quad \varepsilon = \lambda^{-\frac{3}{4p}}, \ \ \frac{3}{4p} < \frac{1}{2}. \tag{6.14}$$

Using $J_0(-z) = J_0(z)$ and the asymptotics (see [1])

$$J_0(z) = \sqrt{\frac{2}{\pi z}}\cos\Big(z - \frac{\pi}{4}\Big) + O\Big(\frac{e^{|\operatorname{Im} z|}}{z^{\frac{3}{2}}}\Big), \qquad |\arg z| \leqslant \frac{\pi}{2},$$

we obtain

$$I_2 = \frac{1}{\lambda^{\frac{1}{4}}}\int_{|t|>\varepsilon}\sqrt{\frac{2}{\pi|t|}}\cos\Big(\sqrt{\lambda}|t| - \frac{\pi}{4}\Big)\,d\nu(t) + \frac{O(1)}{\lambda^{\frac{3}{4}}}\int_{|t|>\varepsilon}\frac{d\nu(t)}{t^{\frac{3}{2}}}, \tag{6.15}$$

where the last term is $O(\lambda^{-\frac{3}{4}})$. Next,

$$|I_1| \leqslant C\int_{|t|<\varepsilon} d\nu(t) \leqslant C\varepsilon^p \int_{|t|<\varepsilon}\frac{d\nu(t)}{|t|^p} = C\gamma\varepsilon^p = O(\lambda^{-\frac{3}{4}}). \tag{6.16}$$

Similarly we have

$$\left|\lambda^{-\frac{1}{4}}\int_{|t|<\varepsilon}\sqrt{\frac{2}{\pi t}}\cos\Big(|t|\sqrt{\lambda} - \frac{\pi}{4}\Big)\,d\nu(t)\right| \leqslant C\lambda^{-\frac{1}{4}}\int_{|t|<\varepsilon}\frac{d\nu(t)}{\sqrt{|t|}} \leqslant C\lambda^{-\frac{1}{4}}\varepsilon^{p-\frac{1}{2}} = O(\lambda^{-\frac{3}{4}}). \tag{6.17}$$

Using (6.13–6.17) and setting $\lambda = \mu_n^0$ gives $\mu_n^1 = \frac{\sigma(\sqrt{\mu_n^0})}{(\mu_n^0)^{\frac{1}{4}}} + O(n^{-\frac{3}{4}})$ which implies (1.3). Moreover, substituting $d\nu(t) = \sum_{k\in\mathbb{Z}}\delta(t - t_k)q_k dt$ into (1.4) we obtain (1.5). ■

7 Appendix

For fixed $\arg\lambda = 2\vartheta$ we have $t = \frac{x}{\sqrt{\lambda}} \in e^{-i\vartheta}\mathbb{R}_+$. We rewrite $t \in S[-\frac{\pi}{2}, 0]$ and ξ in the form

$$t = re^{-i\vartheta} = 1 + \eta e^{-i\varphi}, \qquad \varphi \in [0, \pi], \quad \eta \geqslant 0, \tag{A.1}$$

$$\xi(t) = e^{-i\frac{3\varphi}{2}}\int_0^{\eta}\sqrt{2 + se^{-i\varphi}}\cdot s^{\frac{1}{2}}\,ds, \quad t \in S[-\frac{\pi}{2}, 0]. \tag{A.2}$$

Lemma 7.1. *Let* $t \in S[-\frac{\pi}{2}, 0]$, $R(t) = |\xi(t)|$, $\Phi(t) = \arg \xi(t)$. *Then*

1. $t = re^{-i\vartheta} = 1 + \eta e^{-i\varphi}$, *where* $r, \eta \geqslant 0$, $\vartheta \in [0, \frac{\pi}{2}]$, $\varphi \in [0, \pi]$,
2. $\Phi(t) + \frac{3\varphi}{2} \in \left[-\frac{\vartheta}{2}, 0\right]$,
3. *if* $\eta \leqslant 1$, *then* $\frac{2}{3}|t-1|^{\frac{3}{2}} \leqslant R(t) \leqslant 2|t-1|^{\frac{3}{2}}$,
   if $\eta \geqslant 0$, *then* $\frac{2}{3}\sin^{\frac{3}{2}} \vartheta \leqslant R(t)$,
4. *if* $\vartheta \in (0, \frac{\pi}{2}]$, *then* $\bigcup\limits_{r \in \mathbb{R}_+} \Phi(t) = [-\frac{3\pi}{2}, -2\vartheta)$,
5. *if* $-\pi - \vartheta \leqslant \Phi(t)$, *then* $\arg\left(e^{2i\vartheta}\partial_r \xi(t)\right) \in [-\frac{\pi}{3} + \frac{\vartheta}{6}, \frac{\vartheta}{2}]$.

Proof. The proof of 1 is simple. 2. Consider the integrand in formula (A.2). For $s \in [0, \eta]$ we have $\arg\sqrt{2 + se^{-i\varphi}} \in [-\frac{\vartheta}{2}, 0]$, hence $\Phi(t) + \frac{3\varphi}{2} = \arg\left(\int_0^\eta \sqrt{2 + se^{-i\varphi}} \cdot s^{\frac{1}{2}}\, ds\right) \in [-\frac{\vartheta}{2}, 0]$.

3. Consider the integrand in (A.2). For $s \in [0, \eta]$ we have $1 < \operatorname{Re}\sqrt{2 + se^{-i\varphi}}$ and $|\sqrt{2 + se^{-i\varphi}}| < 3$. Using $R(t) = \left|\int_0^\eta \sqrt{2 + se^{-i\varphi}} \cdot s^{\frac{1}{2}}\, ds\right|$ gives

$$\frac{2}{3}\eta^{\frac{3}{2}} \leqslant R(t) < 3 \cdot \frac{2}{3}\eta^{\frac{3}{2}}, \qquad \eta = |t-1|.$$

The relation $\min\limits_{\arg t = -\vartheta} |t-1| = \sin\vartheta$ for $\vartheta \in [0, \frac{\pi}{2}]$ finishes the proof.

4. Fix $\vartheta \in (0, \frac{\pi}{2}]$. By direct calculation, $\xi(S[-\frac{\pi}{2}, 0]) \subset S[-\frac{3\pi}{2}, 0]$, so $\bigcup\limits_{r \in \mathbb{R}_+} \Phi(t) \subset S[-\frac{3\pi}{2}, 0]$. Using (2.18) we have $\arg\left(e^{2i\vartheta}\xi(t)\right) \to 0$ as $r \to \infty$. Also we have $\xi(0) = i\frac{\pi}{4}$. Since $\xi(t)$ is continuous, this gives $\bigcup_{r \in \mathbb{R}_+} \Phi(t) \supset [-\frac{3\pi}{2}, -2\vartheta)$. Thus we need only prove that $\arg\left(e^{2i\vartheta}\xi(t)\right) < 0$.

Consider $e^{2i\vartheta}\xi(t)$, $t = re^{-i\vartheta}$, as a function of real parameter $r \in \mathbb{R}$. Note that by Lemma 7.2.2 , $\operatorname{Im}\left(e^{2i\vartheta}\xi(t)\right)$ strictly decreases in r. Since $\xi(S[-\frac{\pi}{2}, 0]) \subset S[-\frac{3}{2}\pi, 0]$ and $\xi(0) = i\frac{\pi}{4}$, as r increases $e^{2i\vartheta}\xi(t)$ hits only the negative half of the imaginary axis. Therefore, as soon as $\arg\left(e^{2i\vartheta}\xi(t)\right) > -\frac{\pi}{2}$, we have $\operatorname{Im}\left(e^{2i\vartheta}\xi(t)\right) < 0$. Hence $\arg\left(e^{2i\vartheta}\xi(t)\right) < 0$, which finishes the proof.

5. By (A.2), we have $e^{2i\vartheta}\partial_r\xi(t) = e^{-i\frac{\varphi}{2}} e^{i\vartheta}\sqrt{2\eta}\sqrt{1 + \frac{\eta e^{-i\varphi}}{2}}$, where $\sqrt{1 + \frac{\eta e^{-i\varphi}}{2}} \in S[-\frac{\vartheta}{2}, 0]$. Therefore

$$\arg\left(e^{2i\vartheta}\partial_r\xi(t)\right) \in [-\frac{\varphi}{2} + \frac{\vartheta}{2}, -\frac{\varphi}{2} + \vartheta]. \tag{A.3}$$

Next, by hypothesis, $\Phi \in [-\pi - \theta, -2\vartheta]$. Thus using 2 we obtain $-\varphi \in [-\frac{2}{3}(\pi + \vartheta), -\vartheta]$. Substituting this into (A.3) proves 5. ■

Lemma 7.2. *Let* $t = re^{-i\vartheta} \in S[-\frac{\pi}{2}, 0]$, $R(t) = |\xi(t)|$, $\Phi(t) = \arg\xi(t)$. *Then*

1. $\arg\partial_r\xi(t) \in [-\frac{\pi}{2} - \vartheta, -2\vartheta)$,

2. *if* $\vartheta \in (0, \frac{\pi}{2}]$ *and* $r \geqslant 0$ *then* $\operatorname{Im}\left(e^{2i\vartheta}\partial_r\xi(t)\right) < 0$ *and* $\operatorname{Re}\left(e^{2i\vartheta}\partial_r\xi(t)\right) > 0$,
   if $\vartheta = 0$ *and* $r \in [0,1)$, *then* $\operatorname{Im}\left(e^{2i\vartheta}\partial_r\xi(t)\right) < 0$ *and* $\operatorname{Re}\xi(t) = 0$,
   if $\vartheta = 0$ *and* $r \in (1,\infty)$, *then* $\operatorname{Im}\xi(t) = 0$ *and* $\operatorname{Re}\left(e^{2i\vartheta}\partial_r\xi(t)\right) > 0$,
3. *if* $\Phi(t) \in (-\pi - 2\vartheta, -\frac{\pi}{2} - \vartheta)$, *then* $\partial_r\Phi(t) > 0$,
4. *if* $\Phi(t) \in (-\frac{3\pi}{2} - 2\vartheta, -\pi - \vartheta)$, *then* $\partial_r R(t) < 0$,
5. *if* $\Phi(t) \in (-\frac{\pi}{2} - 2\vartheta, -\vartheta)$, *then* $\partial_r R(t) > 0$.

Proof. 1. By direct calculation, $\partial_r\xi(t) = e^{-i\vartheta}\xi'(t) = e^{-i\vartheta}\sqrt{t^2-1}$. We have $(t^2-1) \in S[-\pi, -2\vartheta)$, so that $\arg\sqrt{|t|^2e^{-2i\vartheta}-1} \in [-\frac{\pi}{2}, -\vartheta)$.

2. For $\vartheta \in (0, \frac{\pi}{2}]$ the result follows from 1. For $\vartheta = 0$, the result follows from (2.17) by direct calculation.

3. We have $\partial_r\Phi(t) = \frac{|\xi'(t)|}{|\xi(t)|}\sin\{\arg\partial_r\xi(t) - \Phi(t)\}$. By 1, $\partial_r\Phi(t)$ is strictly positive for $\Phi(t) \in (-\pi - 2\vartheta, -\frac{\pi}{2} - \vartheta)$.

4. and 5. We have $\partial_r R(t)^2 = 2|\xi'(t)||\xi(t)|\cos\{\arg\partial_r\xi(t) - \Phi(t)\}$. By 1, $\partial_r R(t)$ is positive for $\Phi(t) \in (-\frac{\pi}{2} - 2\vartheta, -\vartheta)$ and negative for $\Phi(t) \in (-\frac{3\pi}{2} - 2\vartheta, -\pi - \vartheta)$. ■

For $\lambda = |\lambda|e^{2i\vartheta} \in \overline{\mathbb{C}}_+ \setminus \{0\}$ we have

$$\frac{2}{3}z(x,\lambda)^{\frac{3}{2}} = \lambda\xi(t), \qquad t = \frac{x}{\sqrt{\lambda}} = re^{-i\vartheta} \in S[-\frac{\pi}{2}, 0]. \tag{A.4}$$

Lemma 7.3. *For each* $\lambda = |\lambda|e^{2i\vartheta} \in S[0,\delta) \setminus \{0\}$ *there exist a unique* $z_* \in \Gamma_\lambda$ *such that* $|z_*| = \min_{z\in\Gamma_\lambda}|z|$. *Moreover, the following relations are valid (*x_* *defined by* $z_* = z(x_*,\lambda)$*):*

1. *if* $\arg\lambda = 0$, *then* $z_* = 0$ *and* $x_* = \sqrt{\lambda}$,
   if $\arg\lambda \in (0,\delta)$, *then* $z_* \in S[-\frac{\pi}{2} - \frac{\vartheta}{2}, -\frac{\pi}{2} + \frac{5}{6}\vartheta]$,
2. $|z(\cdot,\lambda)|$ *is strictly decreasing on* $[0, x_*)$ *and strictly increasing on* (x_*, ∞),
3. *if* $t = \frac{x}{\sqrt{\lambda}}$ *for* $x \in [0, x_*)$, *then* $\arg\left(e^{2i\vartheta}\partial_r\xi(t)\right) \in [-\frac{\pi}{2}, -\frac{\pi}{4} + \frac{3}{4}\vartheta]$,
4. *if* $t_* = \frac{x_*}{\sqrt{\lambda}}$, *then* $|t_*| \leqslant \frac{1}{\sin(\frac{\pi}{3})}$.

Proof. Let us prove uniqueness of z_*. By (A.4), it is sufficient to show that for each λ there exists a unique solution t_* of $\partial_r|\xi(t)| = 0$ for $t = re^{-i\vartheta}$, $r \in \mathbb{R}_+$. For $\arg\lambda = 0$ direct calculation gives $z_* = 0$ and $x_* = \sqrt{\lambda}$. Next we show the existence of t_* in the case $\lambda = |\lambda|e^{2i\vartheta} \in S(0,\delta) \setminus \{0\}$.

Since $\xi(t) \neq 0$, we have $2\partial_r|\xi(t)| = |\xi(t)|^{-1}\operatorname{Re}\left(\xi(t)\overline{\frac{\partial\xi(t)}{\partial r}}\right)$. Let us use the representation (A.1): $t = re^{-i\vartheta} = 1 + \eta e^{-i\varphi}$. For any $t \in S[-\frac{\pi}{2}, 0)$ Lemma 7.1.2 implies $\arg\xi(t) \in$

$[-\frac{3\varphi}{2} - \frac{\vartheta}{2}, -\frac{3\varphi}{2}]$; similarly Lemma 7.2.1 implies $\arg \frac{\partial}{\partial r}\xi(t) \in [-\frac{\varphi}{2} - \frac{3\vartheta}{2}, -\frac{\varphi}{2} - \vartheta]$. Thus $\arg \partial_r |\xi(t)|^2 \in [-\varphi + \frac{\vartheta}{2}, -\varphi + \frac{3\vartheta}{2}]$ and we have

$$\partial_r |\xi(t)| > 0 \qquad\qquad \text{for } -\varphi \in \left(-\frac{\pi}{2} - \frac{\vartheta}{2}, \frac{\pi}{2} - \frac{3\vartheta}{2}\right), \tag{A.5}$$

$$\partial_r |\xi(t)| < 0 \qquad\qquad \text{for } -\varphi \in \left(-\frac{3\pi}{2} - \frac{\vartheta}{2}, -\frac{\pi}{2} - \vartheta\right). \tag{A.6}$$

By (A.5) and (A.6), there exist at least one solution of $\partial_r |\xi(t)| = 0$. Moreover, any solution satisfies

$$-\varphi \in \left[-\frac{\pi}{2} - \frac{3\vartheta}{2}, -\frac{\pi}{2} - \frac{\vartheta}{2}\right]. \tag{A.7}$$

Now we show the uniqueness of t_*. We need only verify that for any t, satisfying (A.7), we have $\frac{1}{2}\frac{\partial^2}{\partial r^2}|\xi(t)|^2 = \left|\frac{\partial \xi(t)}{\partial r}\right|^2 + \mathrm{Re}\left\{\overline{\xi(t)} \cdot \frac{\partial^2 \xi(t)}{\partial r^2}\right\} > 0$. This is true if

$$\left|\partial_r \xi(t)\right|^2 > |\xi(t)| \left|\partial_r^2 \xi(t)\right|. \tag{A.8}$$

We have $\frac{\partial \xi(t)}{\partial r} = e^{-i\vartheta}\sqrt{t^2-1}$, $\frac{\partial^2 \xi(t)}{\partial r^2} = e^{-3i\vartheta}\frac{r}{\sqrt{t^2-1}}$. Therefore (A.8) is equivalent to

$$|(t^2-1)^{\frac{3}{2}}/t| > |\xi(t)|. \tag{A.9}$$

Due to (A.7), $|t| \leqslant 1$. Therefore we have for $t = 1 + \eta e^{-i\varphi}$

$$\frac{|t^2-1|^{\frac{3}{2}}}{|t|} \geqslant \frac{|t-1|^{\frac{3}{2}}|t+1|^{\frac{3}{2}}}{1} = \eta^{\frac{3}{2}}|2 + \eta e^{-i\varphi}|^{\frac{3}{2}}, \qquad |2 + \eta e^{-i\varphi}| \geqslant 2\cos\frac{3\vartheta}{2}. \tag{A.10}$$

Thus using $2\vartheta \in (0, \delta) \subset [0, \frac{4}{3}\arccos\frac{1}{2^{\frac{1}{3}}})$ we obtain

$$\frac{|t^2-1|^{\frac{3}{2}}}{|t|} \geqslant \eta^{\frac{3}{2}}\left(2\cos\frac{3\vartheta}{2}\right)^{\frac{3}{2}} > 2\eta^{\frac{3}{2}}. \tag{A.11}$$

Since $\vartheta \in [0, \delta) \subset [0, \frac{\pi}{3}]$, (A.7) gives $\eta \leqslant 1$. Thus we apply Lemma 7.1.3 , which gives $2\eta^{\frac{3}{2}} \geqslant |\xi(t)|$. Substituting the last estimate in (A.11) yields (A.9) and (A.8). Therefore there exist a unique solution t_* of $\partial_r|\xi(t)| = 0$. Thus $z_* = z(x_*, \lambda)$ for $x_* = t_*\sqrt{\lambda}$.

1. and 2. Using (A.7) and Lemma 7.1.2 we obtain

$$\arg \xi(t_*) \in [-\frac{3\pi}{4} - \frac{11\vartheta}{4}, -\frac{3\pi}{4} - \frac{3\vartheta}{4}]. \tag{A.12}$$

By (A.4), this proves 1. Uniqueness of t_* and the relation (A.4) prove 2.

3. By (A.5), for $r < |t_*|$ we have $-\pi \leqslant -\varphi \leqslant -\frac{\pi}{2} - \frac{\vartheta}{2}$. Using Lemma 7.1.2 we obtain

$$-\frac{\pi}{2} \leqslant \arg\left(e^{2i\vartheta}\partial_r \xi(t)\right) = \arg\left(e^{i\vartheta}e^{-i\frac{\varphi}{2}}\sqrt{1 - \frac{l}{2}e^{-i\varphi}}\right) \leqslant \vartheta - \frac{\varphi}{2} \leqslant -\frac{\pi}{4} + \frac{3\vartheta}{4}.$$

4. Geometric considerations show that in terms of (A.1) $r = \frac{\sin\varphi}{\sin(\varphi - \vartheta)}$. By (A.7) and Lemma 7.1.2 , for $t_* = 1 + \eta_* e^{-i\phi_*}$ we have $\frac{2\pi}{3} - \frac{\vartheta}{3} \geqslant \varphi_* - \vartheta \geqslant \frac{\pi}{3}$, so that $r_* = \frac{\sin\varphi_*}{\sin(\varphi_* - \vartheta)} \leqslant \frac{1}{\sin\frac{\pi}{3}}$.
■

Lemma 7.4. *For each* $\lambda = |\lambda|e^{2i\vartheta} \in S[\delta, \pi] \setminus \{0\}$ *there exist a unique* $z_* \in \Gamma_\lambda$ *such that* $\arg z_* = -\frac{\pi}{3}$. *Moreover, the following relations are valid (here* x_* *defined by* $z_* = z(x_*, \lambda)$*):*

1. *if* $x \in [0, x_*)$, *then* $\arg z(x, \lambda) < -\frac{\pi}{3}$,
   if $x \in (x_*, \infty)$, *then* $\arg z(x, \lambda) > \frac{\pi}{3}$,
2. $|z(\cdot, \lambda)|$ *is strictly increasing on* (x_*, ∞),
3. *if* $t_* = \frac{x_*}{\sqrt{\lambda}}$, *then* $|t_*| \leqslant \frac{1}{\sin\frac{\pi}{3}}$ *uniformly in* λ.

Proof. By Lemma 7.1.4 and (A.4), Γ_λ intersects the ray $\{z : \arg z = -\frac{\pi}{3}\}$ at least once. By (A.4), the sector in z-plane $S[-\frac{\pi}{3} - \varepsilon, -\frac{\pi}{3} + \varepsilon]$ for small $\varepsilon > 0$ corresponds to $S[-\frac{\pi}{2} - 2\vartheta - \frac{3\varepsilon}{2}, -\frac{\pi}{2} - 2\vartheta - \frac{3\varepsilon}{2}]$ in ξ-plane. By Lemma 7.2.3 , in this sector $\partial_r \Phi(t) > 0$. Therefore, by (A.4), $\frac{\partial}{\partial x} \arg z(x, \lambda) > 0$ for $z \in S[-\frac{\pi}{3} - \varepsilon, -\frac{\pi}{3} + \varepsilon]$. Hence z_* is unique and 1 holds.

2. By (A.4), for $x \in (x_*, \infty)$ and $t = \frac{x}{\sqrt{\lambda}}$ the hypothesis of Lemma 7.2.5 is fulfilled. Therefore $\frac{\partial}{\partial r} R(t) > 0$ and $|z(x, \lambda)|$ is strictly increasing in x.

3. By (A.4), we have $\arg \xi(t_*) = -\frac{\pi}{2} - 2\vartheta$. By Lemma 7.1.2 , for $t_* = 1 + \eta_* e^{-i\varphi_*}$ we have $\varphi_* - \vartheta \in [\frac{2\pi}{3} - \frac{\vartheta}{3}, \frac{2\pi}{3}]$. Therefore using the relation $r = \frac{\sin\varphi}{\sin(\varphi - \vartheta)}$ in terms of (A.1), we obtain $|t_*| \leqslant \frac{1}{\sin\frac{\pi}{3}}$. ■

In the next Lemma we analyze the curves $\Gamma_\lambda^\pm$, defined in (2.43).

Lemma 7.5. *Let* $\lambda = |\lambda|e^{2i\vartheta} \in \overline{\mathbb{C}}_+ \setminus \{0\}$. *Then*

1. *if* $\vartheta = 0$, *then* $\Gamma_\lambda = [-\lambda^{\frac{2}{3}}(\frac{3\pi}{8})^{\frac{2}{3}}, \infty)$,
   if $0 < \vartheta \leqslant \frac{\pi}{2}$, *then* $\Gamma_\lambda \subset S[-\pi + \frac{4}{3}\vartheta, 0)$, $\Gamma_\lambda^- \subset S[-\pi + \frac{4}{3}\vartheta, -\frac{\pi}{3}]$, $\Gamma_\lambda^+ \subset S[-\frac{\pi}{2} - \frac{\vartheta}{2}, 0)$,
2. $\inf_{z \in \Gamma_\lambda} |z| \geqslant |\lambda|^{\frac{2}{3}} \sin\vartheta$,
3. $\Gamma_\lambda^- \subset \{z : |z| \leqslant C|\lambda|^{\frac{2}{3}}\}$, *the length of* Γ_λ^- *satisfies* $|\Gamma_\lambda^-| \leqslant C|\lambda|^{\frac{2}{3}}$,
4. *the function* $|z(\cdot, \lambda)|$ *is strictly increasing on* $[x_*, \infty)$.

Proof 1. Direct calculation yields the result for $\vartheta = 0$. For $0 < \vartheta \leqslant \frac{\pi}{2}$ the assertion on Γ_λ follows from Lemma 7.1.4 and $z = \lambda^{\frac{2}{3}}(\frac{3}{2}\xi)^{\frac{2}{3}}$. The assertion on $\Gamma_\lambda^\pm$ for $0 \leqslant \arg\lambda \leqslant \delta$ follows from 1 of the present Lemma and Lemma 7.3.1 . For $\delta \leqslant \arg\lambda \leqslant \pi$ it follows from Lemma 7.4.1 .

2 follows from Lemma 7.1.3 and $z = \lambda^{\frac{2}{3}}(\frac{3}{2}\xi)^{\frac{2}{3}}$.

3. By Lemma 7.3.4 and Lemma 7.4.3 , $|t_*| \leqslant \frac{1}{\sin\frac{\pi}{3}}$ uniformly in $\lambda \in \overline{\mathbb{C}}_+ \setminus \{0\}$. Using the definition (2.19) we conclude that $k(t)$ and $k'(t)$ for $t \in [0, t_*]$ are also uniformly bounded. Now the relations $|\Gamma_\lambda^-| = |\lambda|^{\frac{2}{3}} \int_{[0,t_*]} |k'(t)||dt|$ and $|z(x, \lambda)| = |\lambda|^{\frac{2}{3}}|k(t)|$, $t = \frac{x}{\sqrt{\lambda}}$ complete the proof.

4. We deduce from (A.4) that $\frac{\partial}{\partial x}|z(x,\lambda)| > 0$ if $\partial_r|\xi(t)| > 0$ for $t = re^{-i\vartheta} = \frac{x}{\sqrt{\lambda}}$. Thus for $\delta \leqslant \arg\lambda \leqslant \pi$ the result follows from Lemma 7.4.2 . For $0 \leqslant \arg\lambda \leqslant \delta$ it follows from Lemma 7.3.2 . ■

Proof of Lemma 2.2. Case 1 (or 2) follows from Lemma 7.3 (or Lemma 7.4). ■

Proof of Lemma 2.3. Fix $\lambda = |\lambda|e^{2i\vartheta}$. By (A.4), we have $h(x) = e^{\operatorname{Re}\lambda\xi(t)}$ for $t = \frac{x}{\sqrt{\lambda}} = re^{-i\vartheta}$. By Lemma 7.2.2 , if either $0 < \arg\lambda \leqslant \pi$, $r \geqslant 0$ or $\arg\lambda = 0$, $r > 1$, then $\partial_r \operatorname{Re}(\lambda\xi(t)) > 0$. It remains to consider the case $\vartheta = 0$ and $x \in [0, x_*)$; by Lemma 7.2.2 , $\operatorname{Re}\xi(t) = 0$ and therefore $h = 1$. ■

In order to prove Lemma 2.4, for a fixed $\lambda = |\lambda|e^{2i\vartheta} \in \overline{\mathbb{C}} \setminus \{0\}$ define the function $\Psi(z) = \frac{2}{3}\frac{z^{\frac{3}{2}}}{|\lambda|}$. Ψ maps $\Gamma_\lambda(z_1, z_2)$ onto $\gamma_\lambda(u_1, u_2)$, $u_j = \Psi(z_j)$, $j = 1, 2$. Similarly, we set $\gamma_\lambda(u) = \Psi(\Gamma_\lambda(z))$ for $u = \Psi(z)$ and $\gamma_\lambda^\pm = \Psi(\Gamma_\lambda^\pm)$. We also set $u_0 = \Psi(z_0)$ and $u_* = \Psi(z_*)$.

Consider the integral $\int_{\gamma_\lambda(u)} f(v)|dv|$. By (A.4), we have the parametrization $\gamma_\lambda(u) = \Big\{v \in \mathbb{C} : v = v(r) \quad for \quad r \in [r_u, \infty], \quad \text{where} \quad v(r) = e^{2i\vartheta}\xi(re^{-i\vartheta}), u = \Psi(z(r_u\sqrt{|\lambda|}, \lambda))\Big\}$.

Consider the image of the part of Γ_λ, defined by the hypothesis a) and b) of Lemma 2.4, under the mapping Ψ. Assumption a) transforms into a') $\delta \leqslant \arg\lambda \leqslant \pi$ and $u \in \gamma_\lambda$. Similarly, b) becomes b') $0 \leqslant \arg\lambda \leqslant \delta$, $u \in \gamma_\lambda^+$. Then by Lemma 7.2.2 , $\operatorname{Re} v(r)$ is strictly increasing in r. Thus we choose the variable $\varkappa = \operatorname{Re} v$ and obtain

$$\int_{\gamma_\lambda(u)} |f(v)||dv| = \int_{\operatorname{Re} u}^{\infty} |f(\varkappa + i\operatorname{Im} v(\varkappa))| \left|\frac{dv}{d\varkappa}\right| d\varkappa, \qquad \left|\frac{dv}{d\varkappa}\right| = \sqrt{1 + \left(\frac{d\operatorname{Im} v(\varkappa)}{d\varkappa}\right)^2}.$$

Let us show that in both cases a') and b') the following estimate holds :

$$\int_{\gamma_\lambda(u)} |f(v)||dv| \leqslant C\int_{\operatorname{Re} u}^{\infty} |f(\varkappa + i\operatorname{Im} v(\varkappa))|\, d\varkappa. \tag{A.13}$$

Let us estimate $\left|\frac{dv}{d\varkappa}\right|$. In terms of the parametrization $v(r) = e^{2i\vartheta}\xi(re^{-i\vartheta})$ we have

$$\frac{d\operatorname{Im} v(\varkappa)}{d\varkappa} = \frac{\partial_r \operatorname{Im}\left(e^{2i\vartheta}\xi(t)\right)}{\partial_r \operatorname{Re}\left(e^{2i\vartheta}\xi(t)\right)} = \tan\arg\partial_r e^{2i\vartheta}\xi(t).$$

Consider the two cases a') and b').

a') Let $\delta \leqslant \arg\lambda \leqslant \pi$ and $u \in \gamma_\lambda$. For each point $v(r) \in \gamma_\lambda$ consider $t = re^{-i\vartheta}$ such that $v(r) = e^{2i\vartheta}\xi(t)$. By Lemma 7.2.1 , $\arg\partial_r\xi(t) \in [-\frac{\pi}{2} - \vartheta, -2\vartheta)$. Hence $\arg\partial_r e^{2i\vartheta}\xi(t) \in [-\frac{\pi}{2} + \frac{\delta}{2}, 0)$ and $\left|\frac{d\operatorname{Im} v(\varkappa)}{d\varkappa}\right| \leqslant \frac{1}{\sin(\frac{\delta}{2})}$ uniformly for $\delta \leqslant \arg\lambda \leqslant \pi$. Therefore $\left|\frac{dv}{d\varkappa}\right|$ is also uniformly bounded.

b') Let $0 \leqslant \arg\lambda \leqslant \delta$ and $u \in \gamma_\lambda^+ = \Psi(\Gamma_\lambda^+)$. For each point $v(r) \in \gamma_\lambda^+$ consider $t = re^{-i\vartheta}$ such that $v(r) = e^{2i\vartheta}\xi(t)$. Since $0 \leqslant \vartheta \leqslant \frac{\pi}{7}$, (A.12) (equivalent to Lemma 7.3.1) implies that $-\pi - \vartheta \leqslant \arg\xi(t)$. Therefore by Lemma 7.1.5 , for $v \in \gamma_\lambda^+$ we have $\arg\left(e^{2i\vartheta}\partial_r\xi(t)\right) \in [-\frac{\pi}{3} + \frac{\vartheta}{6}, \frac{\vartheta}{2}]$. Hence $\left|\frac{d\operatorname{Im} v(\varkappa)}{d\varkappa}\right| \leqslant \frac{1}{\sin(\frac{\pi}{3})}$ and $\left|\frac{dv}{d\varkappa}\right|$ is uniformly bounded.

Thus $\left|\frac{dv}{d\varkappa}\right|$ is uniformly bounded in both cases a') and b'), implying (A.13).

Proof of Lemma 2.4. Firstly we prove (2.44) for the case $\delta \leqslant \arg\lambda \leqslant \pi$. By Lemma 7.5.2 , we have $\operatorname{dist}(\Gamma_\lambda, \{0\})$ >const uniformly in λ. Thus we replace $\langle\cdot\rangle$ by $|\cdot|$. The change of variables $v = \Psi(s)$ in (2.44) results in the equivalent relation

$$\int_{\gamma_\lambda(u)} \frac{e^{-2|\lambda|\operatorname{Re} v}}{|v|^{\frac{2}{3}(\alpha+\frac{1}{2})}}|dv| \leqslant \frac{C}{|\lambda|}\frac{e^{-2|\lambda|\operatorname{Re} u}}{|u|^{\frac{2}{3}(\alpha+\frac{1}{2})}}, \quad u = \frac{2}{3}\frac{z^{\frac{3}{2}}}{|\lambda|} \in \gamma_\lambda. \tag{A.14}$$

By (A.13), we have the auxiliary estimate

$$\int_{\gamma_\lambda(u)} e^{-2|\lambda|\operatorname{Re} v}\,|dv| \leqslant C\frac{e^{-2|\lambda|\operatorname{Re} u}}{|\lambda|}, \qquad u\in\gamma_\lambda. \tag{A.15}$$

We show that (A.14) follows from (A.15). If $u\in\gamma_\lambda^+$, then Lemma 7.5.4 yields $|u| = \min\limits_{v\in\gamma_\lambda(u)}|v|$. Therefore (A.14) follows from (A.15). If $u\in\gamma_\lambda^-$, then $\operatorname{dist}(\gamma_\lambda,\{0\})$ >const and we have

$$\int_{\gamma_\lambda(u)} \frac{e^{-2|\lambda|\operatorname{Re} v}}{|v|^{\frac{2}{3}(\alpha+\frac12)}}\,|dv| \leqslant C\int_{\gamma_\lambda(u)} e^{-2|\lambda|\operatorname{Re} v}\,|dv|, \qquad u\in\gamma_\lambda^-. \tag{A.16}$$

By Lemma 7.5.3 , $\gamma_\lambda^-(u)$ is uniformly bounded. Therefore, $|u|$ is bounded on γ_λ^-, so (A.16) and (A.15) imply (A.14).

b) We prove (2.44) for the case $z\in\Gamma_\lambda^+$. It suffices to show that for any $z\in\Gamma_\lambda^+$ we have

$$\int_{\Gamma_\lambda(z)} |e^{-\frac43 s^{\frac32}}|\,|ds| \leqslant C|e^{-\frac43 z^{\frac32}}|, \quad |z|\le 1; \quad \int_{\Gamma_\lambda(z)} \frac{|e^{-\frac43 s^{\frac32}}|}{|s|^{\alpha}}\,|ds| \leqslant C\frac{|e^{-\frac43 z^{\frac32}}|}{|z|^{\alpha+\frac12}}, |z|>1. \tag{A.17}$$

By the change of variable $u=\Psi(z)$ the first estimate in (A.17) follows from (A.15). For the proof of the second estimate in (A.17) we observe that by Lemma 7.5.4 , for $z\in\Gamma_\lambda^+$ we have $|z| = \min\limits_{v\in\Gamma_\lambda(z)}|v|$. Hence, (A.17) follows from (A.15). This proves (2.44).

Next we shall prove (2.45). By the change of variable $u=\Psi(z)$ we have

$$\int_{\Gamma_\lambda(z)} \frac{|ds|}{\langle s\rangle^{\alpha}} = \left(\frac23\right)^{\frac23(\alpha+\frac12)} \frac{I}{|\lambda|^{\frac23(\alpha-1)}}, \quad I=\int_{\gamma_\lambda(u)} \frac{|dv|}{|v|^{\frac13}(\varepsilon+|v|^{\frac23})^{\alpha}}, \quad u=\frac23\frac{z^{\frac32}}{|\lambda|}, \tag{A.18}$$

where $\varepsilon = (\frac32|\lambda|)^{-\frac23}\leqslant C$.

Assume $z\in\Gamma_\lambda^+$, $0<\arg\lambda\leqslant\pi$. Set $a=\operatorname{Re} u$ and $b=\operatorname{Im} u$. By Lemma 7.3.1 , Lemma 7.3.3 , Lemma 7.1.5 and definition of z_*, $b\leqslant 0$. By Lemma 7.2.2 , $\operatorname{Im} v$ is strictly decreasing on γ_λ^+, so that $|b| = \min\limits_{v\in\gamma_\lambda(u)}|\operatorname{Im} v|$. Thus using (A.13) we obtain

$$I\leqslant CJ, \qquad J=\int_a^\infty \frac{d\varkappa}{|\varkappa|^{\frac13}(\varepsilon+|b|^{\frac23}+|\varkappa|^{\frac23})^{\alpha}}.$$

Consider two cases: $a\geqslant 0$ and $a<0$. Firstly, let $a\geqslant 0$. Then we have

$$J=\frac32\int_{a^{\frac23}}^\infty \frac{dx}{(\varepsilon+|b|^{\frac23}+x)^{\alpha}} = \frac{\frac32}{\alpha-1}\frac{1}{(\varepsilon+|b|^{\frac23}+a^{\frac23})^{\alpha-1}} \leqslant \frac{C}{(\varepsilon+|u|^{\frac23})^{\alpha-1}}$$

Therefore

$$\frac{I}{|\lambda|^{\frac23(\alpha-1)}} \leqslant \frac{C}{|\lambda|^{\frac23(\alpha-1)}(\varepsilon+|u|^{\frac23})^{\alpha-1}} = \frac{C}{\langle z\rangle^{\alpha}},$$

which together with (A.18) proves (2.45). Secondly, let $a < 0$. Again, by $x = \varkappa^{\frac{2}{3}}$, we obtain

$$J \leqslant C \int_0^\infty \frac{d\varkappa}{\varkappa^{\frac{1}{3}}(\varepsilon + |b|^{\frac{2}{3}} + \varkappa^{\frac{2}{3}})^\alpha} \leqslant \frac{C}{(\varepsilon + |b|^{\frac{2}{3}})^{\alpha-1}}.$$

Due to Lemma 7.3.1 and Lemma 7.4.1 , for $u = a + ib \in \gamma_\lambda^+$ and $a < 0$ we have $|a| \leqslant C|b|$ uniformly in $0 \leqslant \arg\lambda \leqslant \pi$. Therefore $|u| \leqslant Cb$ and $\frac{1}{(\varepsilon+|b|^{\frac{2}{3}})^{\alpha-1}} \leqslant \frac{C}{(\varepsilon+|u|^{\frac{2}{3}})^{\alpha-1}}$. Thus

$$\frac{I}{|\lambda|^{\frac{2}{3}(\alpha-1)}} \leqslant \frac{C}{|\lambda|^{\frac{2}{3}(\alpha-1)}(\varepsilon + |u|^{\frac{2}{3}})^{\alpha-1}} = \frac{C}{\langle z\rangle^\alpha}, \tag{A.19}$$

which together with (A.18) proves (2.45).

Assume $z \in \Gamma_\lambda^-$ and $\delta \leqslant \arg\lambda \leqslant \pi$. Recall that $u_* = \frac{2}{3}\frac{z_*^{\frac{3}{2}}}{|\lambda|}$. We have $I = I_- + I_+$, where

$$I_- = \int_{\gamma_\lambda(u,u_*)} \frac{|dv|}{|v|^{\frac{1}{3}}(\varepsilon + |v|^{\frac{2}{3}})^\alpha}, \quad I_+ = \int_{\gamma_\lambda^+} \frac{|dv|}{|v|^{\frac{1}{3}}(\varepsilon + |v|^{\frac{2}{3}})^\alpha}, \quad u = \frac{2}{3}\frac{z^{\frac{3}{2}}}{|\lambda|}.$$

By Lemma 7.5.2 , we have $|z| \geqslant \sin\frac{\delta}{2}$ for $z \in \Gamma_\lambda$, so $|u| \geqslant \frac{2}{3}\left(\sin\frac{\delta}{2}\right)^{\frac{3}{2}}$ for $u \in \gamma_\lambda$. By Lemma 7.5.3 , the length of the curve $|\gamma_\lambda^-| < C$. Therefore $I_- \leqslant C$. By (A.19) for $u = u_*$, we have $I_+ \leqslant C$ so that $I \leqslant C$. Next, by Lemma 7.5.3 , γ_λ^- is uniformly bounded, implying $C \leqslant (\varepsilon + |u|^{\frac{2}{3}})^{-1\alpha}$ and

$$\frac{I}{|\lambda|^{\frac{2}{3}(\alpha-1)}} \leqslant \frac{C}{|\lambda|^{\frac{2}{3}(\alpha-1)}(\varepsilon + |u|^{\frac{2}{3}})^{\alpha-1}} = \frac{C}{\langle z\rangle^\alpha}, \tag{A.20}$$

which together with (A.18) proves (2.45). ■

Proof of Lemma 2.5. First we prove (2.46). By the change of variable $u = \Psi(z)$, we have

$$\int_{\Gamma_\lambda^-} \frac{|ds|}{\langle s\rangle^\alpha} = \left(\frac{2}{3}\right)^{\frac{2}{3}(\alpha+\frac{1}{2})} \frac{I}{|\lambda|^{\frac{2}{3}(\alpha-1)}}, \quad I = \int_{\gamma_\lambda^-} \frac{|dv|}{|v|^{\frac{1}{3}}(\varepsilon + |v|^{\frac{2}{3}})^\alpha}, \tag{A.21}$$

where $\varepsilon = (\frac{3}{2}|\lambda|)^{-\frac{2}{3}} \leqslant C$. By Lemma 7.2.2 , $\operatorname{Im} v$ is strictly decreasing on γ_λ^-. Thus we parameterize the last integral by $\chi = \operatorname{Im} v$, so that $v(\chi) = \operatorname{Re} v(\chi) + i\chi$. By Lemma 7.5.3 , $\gamma_\lambda^- \subset \{u \in \mathbb{C} : |u| < c\}$ for some $c > 0$ independent of λ. Therefore

$$I \leqslant 2\int_0^c \left|\frac{dv}{d\chi}\right| \frac{d\chi}{\chi^{\frac{1}{3}}\{\varepsilon + \chi^{\frac{2}{3}}\}^\alpha}, \tag{A.22}$$

where $\left|\frac{dv}{d\chi}\right| = \sqrt{1 + \left(\frac{d\operatorname{Re} v(\chi)}{d\chi}\right)^2}$. Recall that $\lambda = |\lambda|e^{2i\vartheta}$ and $t = re^{-i\vartheta}$. In order to estimate $\left|\frac{d\operatorname{Re} v(\chi)}{d\chi}\right|$ we make use of the parametrization

$\gamma_\lambda(u)=\Big\{v\in\mathbb{C}: v=v(r)\ for\ r\in[r_u,\infty],\ \text{where } v(r)=e^{2i\vartheta}\xi(re^{-i\vartheta}), u=\Psi(z(r_u\sqrt{|\lambda|},\lambda))\Big\}.$
Thus we have

$$\frac{d\operatorname{Re}v(\chi)}{d\chi}=\frac{\partial_r\operatorname{Re}\left(e^{2i\vartheta}\xi(t)\right)}{\partial_r\operatorname{Im}\left(e^{2i\vartheta}\xi(t)\right)}=\cot\arg\left(e^{2i\vartheta}\partial_r\xi(t)\right).$$

By Lemma 7.3.3 , Lemma 7.4.1 , (2.42) and Lemma 7.1.5 , $e^{2i\vartheta}\partial_r\xi(t)$ is in a sector isolated from the real axis uniformly in λ. Therefore $\left|\frac{d\operatorname{Re}v(\chi)}{d\chi}\right|\leqslant C$ and $\left|\frac{dv}{d\chi}\right|\leqslant C$. Substituting this estimate in (A.21) and making change of variable $x=\varepsilon^{-\frac32}\chi$ we obtain

$$I\leqslant\frac{C}{\varepsilon^{\alpha-1}}\int_0^{C_1\varepsilon^{-\frac32}}\frac{dx}{x^{\frac13}(1+x^{\frac23})^{\alpha}},\qquad \varepsilon=(\frac32|\lambda|)^{-\frac23}\leqslant C,\tag{A.23}$$

Recall that $|\lambda|\geqslant R>0$. Then the proof of (2.46) follows from (A.21) and (A.23).

Now we prove (2.47). By the change of variable $u=\Psi(z)$, we have

$$\int_{\Gamma_\lambda}\frac{|ds|}{|\lambda|^{\frac43}+|s|^2}=\frac{I_-+I_+}{|\lambda|^{\frac23}},\quad I_\pm=\int_{\gamma_\lambda^\pm}\frac{|dv|}{(\frac32|v|)^{\frac13}(1+(\frac32|v|)^{\frac43})},\tag{A.24}$$

For I_+ we use (A.13), which gives

$$I_+\leqslant C\int_0^\infty\frac{d\varkappa}{\varkappa^{\frac13}(1+\varkappa^{\frac43})}\leqslant C.\tag{A.25}$$

In order to estimate I_- we use the parametrization of γ_λ^- by $\chi=\operatorname{Im}v$. Repeating the arguments used above for the proof of (2.46), we conclude that $\left|\frac{dv}{d\chi}\right|\leqslant C$ and

$$I_-\leqslant C\int_0^\infty\frac{d\chi}{\chi^{\frac13}(1+\chi^{\frac43})}\leqslant C.\tag{A.26}$$

Now (A.25) and (A.26) give $I\leqslant C$, which by (A.24) proves (2.47). ■

Let $P_\pm$ be defined by (4.27).

Lemma 7.6. *Let $\lambda=|\lambda|e^{2i\vartheta}\in S[0,\delta]$ and $q,q'\in L^\infty(\mathbb{R})$. Then for some absolute constant C the following estimates are fulfilled:*

$$|P_-(z,z_*)|\leqslant C|e^{-\frac43 z^{\frac32}}|\cdot\Big(\|q'\|_\infty+\|q\|_\infty|\lambda|^{-\frac16}\Big),\quad z\in\Gamma_\lambda^-,\tag{A.27}$$

$$|P_+(z_1,z_2)|\leqslant C|e^{\frac43 z_2^{\frac32}}|\cdot\Big(\|q'\|_\infty+\|q\|_\infty|\lambda|^{-\frac16}\Big).\tag{A.28}$$

Proof. Let $x_1\leqslant x_2\leqslant x_*$, $t_{1,2}=\frac{x_{1,2}}{\sqrt{\lambda}}$ and $z_{1,2}=z(x_{1,2},\lambda)\in\Gamma_\lambda^-$. Let $\xi_{1,2}=\xi(t_{1,2})$. Then, Lemma 7.3.2 yields $|\xi_1-\xi_2|\leqslant 2|\xi_1|$. Lemma 2.3 gives $|e^{2\lambda(\xi_1-\xi_2)}-1|\leqslant 2|\lambda||\xi_1-\xi_2|$ and $|e^{2\lambda(\xi_1-\xi_2)}-1|\leqslant C$. Therefore

$$|(e^{2\lambda(\xi_1-\xi_2)}-1)/\lambda\xi_1|\leqslant C(1+|\lambda||\xi_1|)^{-1}.\tag{A.29}$$

Using $z_{1,2} = \lambda^{\frac{2}{3}}(\frac{3}{2}\xi_{1,2})^{\frac{2}{3}}$ we obtain the following estimate uniformly in $0 \leqslant \arg\lambda \leqslant \delta$:

$$|(e^{\frac{4}{3}(z_1^{\frac{3}{2}} - z_2^{\frac{3}{2}})} - 1)z^{-\frac{3}{2}}| \leqslant C\langle z_1\rangle^{\frac{3}{2}}. \tag{A.30}$$

Introduce the functions $R_\pm(s,z) = e^{\pm\frac{4}{3}s^{\frac{3}{2}}} - e^{\pm\frac{4}{3}z^{\frac{3}{2}}}$. Evidently $\partial_s R_\pm(s,z) = \pm 2\sqrt{s}e^{\pm\frac{4}{3}s^{\frac{3}{2}}}$. Integrating by parts, we have

$$P_-(z_*, z) = R_-(z, z_*)f(z) + \int_{\Gamma_\lambda(z,z_*)} R_-(s, z_*)f'(s)\,ds, \quad f(s) = \rho(s)\hat{q}(s)/(2\sqrt{s}\lambda^{\frac{1}{6}}). \tag{A.31}$$

Therefore using (2.29), (2.30), (A.30) and $|\frac{d}{ds}\hat{q}(s)| \leqslant C\|q'\|_\infty|\lambda|^{-\frac{1}{6}}$ we obtain

$$|P(z,z_*)| \leqslant C\frac{|e^{-\frac{4}{3}z^{\frac{3}{2}}}|}{|\lambda|^{\frac{1}{6}}}\left[\frac{\|q\|_\infty}{\langle z\rangle^{\frac{1}{2}}} + \int_{\Gamma_\lambda(z,z_*)}|ds|\left(\frac{|\lambda|^{-\frac{1}{6}}\|q'\|_\infty}{\langle s\rangle^{\frac{1}{2}}} + \frac{|\lambda|^{-\frac{2}{3}}\|q\|_\infty}{\langle s\rangle^{\frac{1}{2}}} + \frac{\|q\|_\infty}{\langle s\rangle^{\frac{3}{2}}}\right)\right], \tag{A.32}$$

which together with (2.45–2.46) gives (A.27). Similarly we have

$$P_+(z_1, z_2) = -R_+(z_1, z_2)f(z_1) - \int_{\Gamma_\lambda(z_1,z_2)} R_+(s, z_2)f'(s)ds. \tag{A.33}$$

Again using (2.29), (2.30), (A.30) and $|\frac{d}{ds}\hat{q}(s)| \leqslant C\|q'\|_\infty|\lambda|^{-\frac{1}{6}}$ we have

$$|P(z_1,z_2)| \leqslant C\frac{|e^{\frac{4}{3}z_2^{\frac{3}{2}}}|}{|\lambda|^{\frac{1}{6}}}\left[\frac{\|q\|_\infty}{(1+|z_2|)^{\frac{1}{2}}} + \int\limits_{\Gamma_\lambda(z_1,z_2)}\left(\frac{|\lambda|^{-\frac{1}{6}}\|q'\|_\infty}{\langle s\rangle^{\frac{1}{2}}} + \frac{|\lambda|^{-\frac{2}{3}}\|q\|_\infty}{\langle s\rangle^{\frac{1}{2}}} + \frac{\|q\|_\infty}{\langle s\rangle^{\frac{3}{2}}}\right)|ds|\right],$$

which together with (2.45), (2.46) gives (A.28). ■

Lemma 7.7. *Let* $q \in \mathcal{B}$. *Define* $F(z) = a^2(ze^{\pm\frac{2\pi i}{3}})V_q(z)$. *Then uniformly in* $z \in \Gamma_\lambda^-$ *and* $\lambda \in S[-\delta,\delta]$ *the following estimates hold for sufficiently large* $|\lambda|$*:*

$$\left|\int_{\Gamma_\lambda(z)} e^{\frac{4}{3}(z^{\frac{3}{2}} - s^{\frac{3}{2}})}F(s)\,ds\right| \leqslant C\frac{\|q\|_{\mathcal{B}}}{|\lambda|^{\frac{1}{3}}}, \qquad \left|\int_{\Gamma_\lambda(z,z_*)} e^{\frac{4}{3}s^{\frac{3}{2}}}F(s)\,ds\right| \leqslant C\frac{\|q\|_{\mathcal{B}}}{|\lambda|^{\frac{1}{3}}}. \tag{A.34}$$

Proof. We show the first estimate in (A.34).Using (2.29), (2.30), (2.38) and (2.44) we obtain

$$\left|\int_{\Gamma_\lambda^+} e^{\frac{4}{3}(z^{\frac{3}{2}} - s^{\frac{3}{2}})}F(s)\,ds\right| \leqslant C\cdot\frac{\|q\|_\infty}{|\lambda|^{\frac{1}{3}}}. \tag{A.35}$$

Using $R_-(s,z_*) = e^{-\frac{4}{3}s^{\frac{3}{2}}} - e^{-\frac{4}{3}z_*^{\frac{3}{2}}}$ we integrate by parts the integral over $\Gamma_\lambda(z,z_*)$. We have

$$I = \int_{\Gamma_\lambda(z,z_*)} e^{\frac{4}{3}(z^{\frac{3}{2}} - s^{\frac{3}{2}})}F(s)ds = \frac{e^{\frac{4}{3}z^{\frac{3}{2}}}}{2}\left[R_-(z,z_*)\frac{F(z)}{\sqrt{z}} + \int_{\Gamma_\lambda(z,z_*)} R_-(s,z_*)\left(\frac{F(s)}{\sqrt{s}}\right)'\,ds\right]. \tag{A.36}$$

Therefore, using Lemma 2.3, $|\frac{d}{ds}\hat{q}(s)| \leqslant C|\lambda|^{-\frac{1}{6}}\|q'\|_\infty$, (2.29), (2.30), (2.38), (2.39) and (A.30), we obtain from (A.36)

$$|I| \leqslant \frac{C}{|\lambda|^{\frac{1}{3}}}\left[\frac{\|q\|_\infty}{\langle z\rangle^{\frac{1}{2}}} + \int_{\Gamma_\lambda^-}\left(\frac{\|q\|_\infty}{\langle s\rangle^{\frac{3}{2}+\frac{1}{2}}} + \frac{|\lambda|^{-\frac{1}{6}}\|q'\|_\infty}{\langle s\rangle} + \frac{|\lambda|^{-\frac{2}{3}}\|q\|_\infty}{\langle s\rangle}\right)|ds|\right]. \tag{A.37}$$

By Lemma 2.5, for sufficiently large $|\lambda|$ this implies $|I| \leqslant C|\lambda|^{-\frac{1}{3}}\|q\|_{\mathcal{B}}\Big(1 + |\lambda|^{-\frac{1}{6}}\log(|\lambda|+1)\Big)$. Together with (A.35) this proves the first estimate in (A.34). In order to prove the second estimate in (A.34) we use similar arguments with $R_+(s, z_*) = e^{\frac{4}{3}s^{\frac{3}{2}}} - e^{\frac{4}{3}z_*^{\frac{3}{2}}}$. ■

References

[1] Abramowitz, M. and Stegun, A., eds.: *Handbook of Mathematical Functions.* N.Y.: Dover Publications Inc.

[2] Chelkak, D., Kargaev, P., Korotyaev, E.: An Inverse Problem for an Harmonic Oscillator Perturbed by Potential: Uniqueness. Lett. Math. Phys. **64**(1), 7-21 (2003)

[3] Chelkak D., Kargaev P., Korotyaev E.: Inverse problem for harmonic oscillator perturbed by potential, characterization. Comm. Math. Phys. in press

[4] Pöschel, J., Trubovitz, E.: *Inverse Spectral Theory.* Boston: Academic Press, 1987